

4947
v. 73
#6

PHYTOLOGIA

An international journal to expedite plant systematic, phytogeographical
and ecological publication

Vol. 73

December 1992

No. 6

CONTENTS

NESOM, G.L., A new species of <i>Sisyrinchium</i> subg. <i>Sisyrinchium</i> (Iridaceae) from México.	429
FERNANDEZ NAVA, R. & N. WAKSMAN, Una especie nueva de <i>Karwinskia</i> (Rhamnaceae) de Tehuacán, Puebla, México.	435
MACROBERTS, M.H. & B.R. MACROBERTS, Observations on toothache grass (<i>Ctenium aromaticum</i> [Poaceae: Chlorideae]) with particular reference to fire.	439
RAYNER, T.G.J., Studies on the genus <i>Bidens</i> L. (Compositae) from the Eastern Hemisphere. 4. A new species from Ethiopia.	445
SEMPLE, J.C., The goldenasters of California, <i>Heterotheca</i> (Compositae: Astereae): New names and combinations.	449
HOCH, P.C. & P.H. RAVEN, <i>Boisduvalia</i> , a coma-less <i>Epilobium</i> (Onagraceae).	456
HOCH, P.C., A new species of <i>Epilobium</i> (Onagraceae) in California. ..	460
Erratum.	463
Books received.	464
Index to authors, volume 73.	473
Index to taxa, volume 73.	474

Contents continued on the inside cover.

LIBRARY

FEB - 8 1993

NEW YORK
BOTANICAL GARDEN

(Contents continued)

Corrections and additions.	509
Index to reviewers, volume 73.	510-511
Disposition of manuscripts received for volumes 72 and 73.	512

PHYTOLOGIA (ISSN 00319430) is published monthly with two volumes per year by Michael J. Warnock, 185 Westridge Drive, Huntsville, TX 77340. Second Class postage at Huntsville, TX. Copyright ©1991 by PHYTOLOGIA. Annual domestic individual subscription (12 issues): \$36.00. Annual domestic institutional subscription (12 issues): \$40.00. Foreign and/or airmail postage extra. Single copy sales: Current issue and back issues volume 67 to present, \$3.50; Back issues (previous to volume 67), \$3.00 (add \$.50 per copy postage and handling US [\$1.00 per copy foreign]). Back issue sales by volume: \$17.00 per volume 42-66 (not all available as complete volumes); \$21.00 per volume 67-present; add \$2.00 per volume postage US (\$4.00 per volume foreign). POSTMASTER: Send address changes to Phytologia, 185 Westridge Drive, Huntsville, TX 77340.

A NEW SPECIES OF *SISYRINCHIUM* SUBG. *SISYRINCHIUM* (IRIDACEAE) FROM MEXICO

Guy L. Nesom & Luis Hernández S.

Department of Botany, University of Texas, Austin, Texas 78713 U.S.A.

ABSTRACT

Sisyrinchium novoleonense is described from Nuevo León and Coahuila, México, where it is abundant in areas of relatively high elevation around Cerro Potosí, Peña Nevada, and Sierra de Arteaga. It is a member of subg. *Sisyrinchium* and apparently related to the *S. scabrum* species group, but it clearly stands apart from these plants in its long rhizomes, large flowers with a long filament tube, and habitats at higher elevations. Based on records in LL, TEX, five other species of subg. *Sisyrinchium* are known from Coahuila, Nuevo León, and Tamaulipas: *S. biforme*, *S. micranthum*, *S. dimorphum*, *S. demissum*, and *S. scabrum*. The latter two probably are the closest relatives of *S. novoleonense* but they are more similar between themselves than is either to *S. novoleonense*.

KEY WORDS: *Sisyrinchium*, Iridaceae, México

RESUMEN

Se describe *Sisyrinchium novoleonense* de Nuevo León y Coahuila, México, abundante en áreas que sobrepasan los 2700 metros de altitud del Cerro Potosí, Peña Nevada, y Sierra de Arteaga. La nueva especie pertenece al subg. *Sisyrinchium* y aparentemente esta relacionada con el grupo de especies de *S. scabrum*, sin embargo difiere claramente de esta por el mayor tamaño de rizoma, flor, y tubo estaminal y su habitat en elevaciones mayores. Con base en registros en LL, TEX, se conocen cinco especies mas del subg. *Sisyrinchium* de Coahuila, Nuevo León, y Tamaulipas: *S. biforme*, *S. micranthum*, *S. dimorphum*, *S. demissum*, y *S. scabrum*. Aunque los dos últimos son probablemente los mas cercanos a *S. novoleonense*, presentan mayor similitud entre sí que cualquiera de ellos con *S. novoleonense*.

PALABRAS CLAVE: *Sisyrinchium*, Iridaceae, México

Recent studies have contributed greatly to an understanding of the phylogeny and taxonomy of *Sisyrinchium* L. and its close relatives (Rudall *et al.* 1986; Goldblatt *et al.* 1989 and 1990; Goldblatt 1990). *Sisyrinchium* is one of the largest genera within the tribe Sisyrinchieae Baker as well as the whole family, and the genus is well known for seemingly small and variable differences among some of its species. Each of four relatively recent papers (Oliver 1969; Calderón & Rzedowski 1985; McVaugh 1989; Henrich & Goldblatt 1987) has described a new species of *Sisyrinchium* from México, and there are now ca. 32 Mexican species (depending on concepts of species delimitation), seven in subg. *Sisyrinchium* and 25 in subg. *Echthronema* (Herbert) Goldbl. (sometimes treated as the genus *Hydastylus* Salisb.). Three recent treatments have clarified the taxonomy of the genus in the southern half of México, where more than two-thirds of the Mexican species occur (Henrich & Goldblatt 1987; McVaugh 1989; Calderón & Rzedowski 1990). In the course of an attempt to identify the *Sisyrinchium* of northeastern México, particularly in relation to several floristic surveys in progress within the area, it has become evident that the following species of sect. *Sisyrinchium* is undescribed.

Sisyrinchium novoleonense Nesom & L. Hernández S., *sp. nov.* TYPE: MEXICO. Nuevo León: Mpio. Dr. Arroyo, Peña Nevada area, ca. 35 km ENE of Dr. Arroyo, NW slope of Picacho Onofre, 28 Jul 1977, C. Wells & G. Nesom 394 (HOLOTYPE: LL!).

Sisyrinchio scabro Cham. & Schlecht. similis sed rhizomatibus valde evolutis, floribus paucioribus in quaque spatha, tepalis majoribus, et tubo staminali longiore differt.

Perennials from a short (5-20 mm) rhizome bearing fibrous roots up to 10 cm long, thickened but not fleshy or swollen, slightly tapering distally; herbage glabrous but the leaves and stems usually beset with minute marginal papillae to hairlike teeth. Stems 1-2 per plant, erect, 2-4 dm high, compressed and 2 winged, 2-3 mm wide, slightly bent at the nodes, with 1-2 branch nodes on the upper 1/2-1/3, each node with a leaflike bract and a terminal branch plus 1-2 axillary branches, all branches of equal to subequal length, each bearing a spathe at the apex. Leaves linear, becoming linear-lanceolate near the stem apex, 2-3 mm wide, the basal 7-20 cm long, the cauline (nodal bracts) like the basal but smaller, 5-10 cm long, leaves absent below the nodes. Spathes 1-2(-4) flowered, compressed, narrowly lanceolate-elliptic in outline, 18-25 mm long, 2.5-3.0 mm wide, the bracts ovate, equal to subequal, smooth margined, the outer bract fused 3-6 mm above the base, commonly with a broad hyaline

margin near the base; pedicels erect to slightly spreading in flower and fruit, about the same length as the spathe. Ovary obpyriform, minutely pilose with delicate, glandular hairs; tepals blue to bluish purple, obovate-elliptic, 15-18 mm long, apically rounded to slightly retuse, with an aristate extension up to 1.5 mm long; filaments completely united into a filament tube 6-7 mm long, anthers erect, verticillate and essentially sessile at the summit of the column, 0.8-1.0 mm long, the style branches not or scarcely projecting beyond the tips of the anthers. Fruits ca. 4 mm long, 4-5 mm wide, blocky-globose, truncate at the apex, 3 angled and weakly 3 sulcate; seeds black, obscurely pitted, ca. 1 mm in diameter, with a large depression on one side.

Additional collections examined: MEXICO. Coahuila: Mpio. Arteaga, Las Vigas, Cañon de la Carbonera, Sierra de Arteaga, woods of pine, Douglas fir, fir, oak, and *Ceanothus*, 2100-2600 m, 15 Sep 1988, Villarreal et al. 4574 (ASU). Nuevo León: Mpios. Dr. Arroyo and Zaragoza, Peña Nevada area: W side of Picacho Onofre, 3230 m, 4 Jul 1959, Beaman 2692 (TEX); Puerto Mesa del Charco, 3100 m, glade in pine-fir forest, 25 Jun 1978, Hinton et al. 17368 (TEX); NW slope of Picacho Onofre, 30 Jul 1977, Wells & Nesom 448 (LL). Mpio. Galeana, Cerro Potosí: SE side, above Ejido 18 Marzo, open pine forest, ca. 3020 m, 25 Jun 1960, Beaman 3315 (TEX); lower slopes, pine woods, 6-8000 ft, 27 Aug 1987, Bogler & Atkins 203 (TEX); 8 mi up from 18 de Marzo, pine-oak transition, 24 Aug 1984, Lavin 4815 (TEX); near microwave tower, 9000 ft, 7 Jul 1963, McGregor et al. 259 (LL); clearing in pine forest, 3250 m, 8 Aug 1970, Hinton et al. 17298 (TEX); NE face, oak forest, ca. 2800 m, 24 Jul 1977, Wells & Nesom 226 (LL): Cerro Potosí area, Mpio. Galeana: San José Las Joyas (N slope of CP), open pine forest, 2730 m, 28 Aug 1983, Hinton et al. 18584 (TEX); ascent to Sierra Infernillo, ca. 15 mi SW of Galeana, 9-10000 ft, pine savanna, 16 Jun 1934, Mueller 833 (TEX).

Sisyrinchium novoleonense is clearly a member of subg. *Sisyrinchium* in its blue tepals and sessile stamens borne at the apex of a filament tube. Specimens of the new species have accumulated at LL, TEX under the name of *S. quadrangulatum* Klatt, which is superficially similar in habit, but the latter is a species restricted to central México in the high mountains from Pico de Orizaba (the type locality) in Veracruz and Puebla to Nevado de Toluca in Edo. México (see Rzedowski & Rzedowski 1985). There has been some confusion about the identity of *S. quadrangulatum* (based on variability in dried flower color, Greenman 1903), but as identified and treated by Calderón & Rzedowski (1990), it is a member of subg. *Echthronema* and not at all related to *S. novoleonense*.

Sisyrinchium novoleonense occurs in the areas of Peña Nevada and Cerro Potosí of Nuevo León, and in the Sierra Arteaga of Coahuila, where it occurs mostly at elevations of 2600-3250 meters. It is known from only one collection in Coahuila (slightly north of Cerro Potosí), but it should be expected to occur more abundantly there, as do many other species with a similar pattern

of geographic distribution.

Based on records of specimens in LL,TEX, ca. 11 species of *Sisyrinchium* (acknowledging the caveats below regarding taxonomic difficulties within the *S. scabrum* Schlecht. & Cham. group) in two subgenera are known to occur in the northeastern Mexican states of Coahuila, Nuevo León, and Tamaulipas. In subg. *Sisyrinchium* are the following: *S. novoleonense*, the *S. scabrum* group (*S. scabrum*, *S. dimorphum* R. Oliver, and *S. demissum* E. Greene), *S. micranthum* Cav., and *S. biforme* Bickn. *Sisyrinchium biforme* is a species known from coastal sites in Texas and Tamaulipas, not approaching the more montane habitats of the others. Five species of subg. *Echthronema* that occur in Coahuila, Nuevo León, and Tamaulipas are discussed in a separate paper (Nesom in prep.).

Among Mexican and Texan species of subg. *Sisyrinchium*, the new species is similar to *S. scabrum* in its branched stems, distinctively scabrous leaf and stem margins, and minutely pubescent ovaries. It differs from all other Mexican and Texan plants of subg. *Sisyrinchium* particularly in its large flowers (tepals) and long filament tube. It also differs from *S. scabrum* in its fewer flowers per spathe (mostly 1-2 vs. mostly 2-6). In these features, and with its localized geographic distribution and distinctive habitats, it can be identified without equivocation.

Sisyrinchium scabrum has not been recorded for Texas, but Oliver (1970) noted that *S. ensigerum* Bickn. (the type from Bexar Co., Texas) shows similarities to it in trans-Pecos Texas, and the two probably are conspecific. Both also are similar to *S. dimorphum* (the type from Valverde Co., Texas), which Oliver (1970) distinguished from *S. ensigerum* in its relatively longer stems, spreading-deflexing pedicels shorter than the spathes at anthesis, and its olivaceous color when dried; also, the ovaries of *S. dimorphum* are glabrous, compared to the minutely hairy ones of *S. scabrum* and other close relatives. Plants of both species are very similar in vegetative and floral morphology, lacking well developed rhizomes and with tepals mostly 7-9 mm long and a filament tube 2-3(-4) mm long. Although *S. dimorphum* is primarily endemic to the Edwards Plateau of Texas, it is also known from a few scattered localities in northern Coahuila, Nuevo León, and Tamaulipas. Plants in Chiapas and Guatemala identified as *S. dimorphum* (Henrich & Goldblatt 1987) are more similar in both habit and habitat to *S. biforme* Bickn. and are better identified as the latter, although this needs further investigation. *Sisyrinchium scabrum* is widely distributed in México, from Oaxaca to Veracruz, Nuevo León, and Coahuila and to Jalisco and Chihuahua in the western part of the country. The latter is abundant in Nuevo León where it occurs at relatively low elevations (ca. 1500-2700 m) compared to *S. novoleonense*. There are numerous specimens in LL,TEX of *S. scabrum* collected from the foothills of both Peña Nevada and Cerro Potosí.

Sisyrinchium micranthum Cav., another species of subg. *Sisyrinchium* pe-

ripherally related to *S. scabrum*, is known from a few collections in south-central Nuevo León at the northern limit of its distribution, extending from there southward through San Luis Potosí and Veracruz into Central and South America. It sometimes may appear similar to *S. scabrum*, but plants of *S. micranthum* are annual, without any rhizome, and they generally produce shorter stems with 0-1 branches, reflexing pedicels, mostly glabrous ovaries, and smaller spathes, tepals, anthers, and capsules.

Also closely related to *Sisyrinchium scabrum* is what can be identified in northern Coahuila as *S. demissum* (the type from Arizona). It differs from *S. scabrum* and *S. dimorphum* in its more strongly developed rhizome and erect pedicels that conspicuously exceed the spathe. *Sisyrinchium demissum*, however, produces flowers in the same size range of *S. scabrum* and *S. dimorphum* and appears to intergrade with *S. scabrum*. Further northwest, formal varieties have been recognized within *S. demissum* (Kearney & Peebles 1960), which appears to intergrade with yet other species (Holmgren 1977; Cholewa & Henderson 1984). A more detailed study is needed of the *S. scabrum* group, particularly in northern México and the southcentral United States, before an accurate accounting can be given of the variation patterns and corresponding nomenclature.

ACKNOWLEDGMENTS

We thank Billie Turner and Carol Todzia for their review of the manuscript and the staff of ASU for a loan of specimens.

LITERATURE CITED

- Calderón de Rzedowski, G. & J. Rzedowski. 1985. *Sisyrinchium conzattii* (Iridaceae), una nueva especie de las altas montañas del centro de México. *Phytologia* 58:371-376.
- . 1990. Iridaceae. Pp. 323-340 in *Flora Fanerogámica del Valle de México*. Vol. III. Monocotyledoneae. Inst. Ecología, Pátzcuaro, Michoacán, México.
- Cholewa, A.F. & D.M. Henderson. 1984. Biosystematics of *Sisyrinchium* section *Bermudiana* (Iridaceae) of the Rocky Mountains. *Brittonia* 36:342-363.
- Goldblatt, P. 1990. Phylogeny and classification of Iridaceae. *Ann. Missouri Bot. Gard.* 77:607-627.

- Goldblatt, P., J.E. Henrich, & R.C. Keating. 1989. Seed morphology of *Sisyrinchium* (Iridaceae - Sisyrinchieae) and its allies. Ann. Missouri Bot. Gard. 76:1109-1117.
- Goldblatt, P., P. Rudall, & J.E. Henrich. 1990. The genera of the *Sisyrinchium* alliance (Iridaceae: Iridoideae): phylogeny and relationships. Syst. Bot. 15:497-510.
- Greenman, J. 1903. New and otherwise noteworthy angiosperms from México and Central America. Proc. Amer. Acad. Arts 39:69-120.
- Henrich, J.E. & P. Goldblatt. 1987. Mesoamerican *Sisyrinchium* (Iridaceae): New species and records and notes on typification. Ann. Missouri Bot. Gard. 74:903-910.
- Holmgren, P.K. 1977. *Sisyrinchium*. Intermountain Fl. 6:538-543.
- Kearney, T.H. & R.H. Peebles. 1960. *Arizona Flora* (ed. 2). University of California Press, Berkeley, California.
- McVaugh, R. 1989. *Sisyrinchium*. Flora Novo-Galiciana 15:311-330.
- Nesom, G.L. In prep. New species of *Sisyrinchium* subg. *Echthronema* (Iridaceae) from northern Mexico.
- Oliver, R.L. 1969. *Sisyrinchium dimorphum* (Iridaceae), a new species from Texas and Mexico. Ann. Missouri Bot. Gard. 55:397.
- . 1970. *Sisyrinchium*. Pp. 425-428 in Correll, D.S. & M.C. Johnston, *Manual of the Vascular Plants of Texas*. Texas Research Foundation, Renner, Texas.
- Rudall, P., A.Y. Kenton, & T.J. Lawrence. 1986. An anatomical and chromosomal investigation of *Sisyrinchium* and allied genera. Bot. Gaz. 147:466-477.

UNA ESPECIE NUEVA DE *KARWINSKIA* (RHAMNACEAE) DE TEHUACAN,
PUEBLA, MEXICO¹

Rafael Fernández Nava

Laboratorio de Botánica Fanerogámica, Escuela Nacional de Ciencias
Biológicas, Instituto Politécnico Nacional, Apartado Postal 17-564, 11410
México, D.F. MEXICO

&

Noemi Waksman

Departamento de Farmacología y Toxicología, Facultad de Medicina,
Universidad Autónoma de Nuevo León, Apartado Postal 146, Col. Del Valle,
Nuevo León, MEXICO

RESUMEN

Se describe *Karwinskia tehuacana* *sp. nov.* planta endémica, del matorral xerófilo que prosperan en la región de Tehuacán, Puebla. La especie parece tener afinidad con *K. humboldtiana* Zucc.

PALABRAS CLAVE: *Karwinskia*, Rhamnaceae, México, especie nueva

ABSTRACT

Karwinskia tehuacana *sp. nov.* is described from the arid scrub of the region of Tehuacán, Puebla. It is related to *K. humboldtiana* Zucc.

KEY WORDS: *Karwinskia*, Rhamnaceae, México, new species

¹Trabajo parcialmente subsidiado por la S.E.P. 88-01-03.

En el año de 1988, se colectó una planta del género *Karwinskia* proveniente de Tehuacán, Puebla y al revisar ejemplares depositados en los herbarios de la Escuela Nacional de Ciencias Biológicas (ENCB) y el Herbario Nacional de México (MEXU) provenientes de esta misma zona se observaron ejemplares pertenecientes al mismo taxa que no se pudieron identificar con la ayuda de la literatura ya existente. Al investigar la identidad de esta especie nos percatamos que se trata de una especie nueva para la ciencia y se describe a continuación.

Karwinskia tehuacana Fernández et Waksman, *sp. nov.* TIPO: MEXICO. Puebla: 6 Km al S de Tepeyehualco, 1 Km al W sobre brecha a la Cascada de Acatzitzimitla, municipio de Atoyatempan, cañada con vegetación de matorral xerófilo, sobre suelo calizo, alt. 1850 m, 28-X-1988, *R. Fernández N. 4975* (HOLOTIPO: ENCB; Duplicados por distribuirse).

Arbor vel frutex, 3-8 m altus; folia simplicia, opposita, petioli 3-5 mm longi, limbi ovato elliptici vel ovato lanceolati, 3-5 cm longi, 1.5-2.0 cm lati, margine integri, supra viridis citrinus, glabratus, infra cum nervis elevatus. Cymae umbelliformes axillares 3-4 florum. Drupae globosae 10-12 mm longae.

Arbol o arbusto de 3-8 m de alto, ramas glabras, grisáceas. Hojas simples opuestas, láminas ovado-elípticas a ovado-lanceoladas de 3-5 cm de largo, 1.5-2.0 cm de ancho, haz verde-limón, glabro, el envés ligeramente más pálido y con las nervaduras secundarias muy prominentes, glabro, margen entero, ápice agudo, base aguda a ligeramente redondeada, pecíolos de 3-5 mm de largo, glabros, estípulas deltoides, ca. 0.5 mm de largo, glabras, persistentes. Inflorescencias dispuestas en cimas axilares de 3-4, pedúnculo de 2-3 mm de largo, glabro; copa floral campanulada de ca. 1.5 mm de largo, glabra; sépalos deltoides de 1.0-1.5 mm de largo, 1.2-1.5 mm de ancho, con abundantes motas negras sobre la superficie; pétalos blanco-amarillentos, unguiculados y cimbiformes de 1.3-1.5 mm de largo, 0.8-1. mm de ancho; estambres envueltos parcialmente por los pétalos, anteras café-amarillentas de 0.5-0.8 mm de largo; disco no muy evidente, verdoso, estilo bilobado en el ápice, estigmas dos papilosos. Fruto drupáceo, verde en la juventud, que se torna a rojo y después a negro en la madurez, globoso a subgloboso de 10-12 mm de diámetro. Semillas 2-3 elipsoides de ca. 5 mm de largo, blanco-amarillentas, lisas.

Material adicional examinado: MEXICO. Puebla: Cascadas de Acatzitzimitla, 9 Km al SE de Tepeyehualco, municipio de Atoyatempan, cañada con vegetación de matorral xerófilo, suelos calizos, alt. 1850 m, 17-XI-1984, *R. Fernández N. 2690* (ENCB); cañada de Acatzitzimitla, municipio de Atoyatempan, cañada con vegetación de matorral alto con *Bambusa*, 28-VI-1982,



F.G. Medrano 12663 (MEXU); 1 Km al W de Nopala, brecha a Atexcal, municipio de Teontepec, cañada con vegetación de matorral calcicola mixto con *Brahea nitida*, *B. dulcis*, *Dasyllirion* sp., 27-IX-1984, *P. Tenorio L. 7455* (ENCB); barranca de Tentzo, 12 Km al W de Molcaxac, carr. a Huatlatlauca, municipio de Huatlatlauca, vegetación de matorral calcicola mixto primario, 22-X-1986, *P. Tenorio L. 12182* (ENCB,MEXU); meseta de San Lorenzo, 8 Km al W de Tehuacán, camino a Tecamachalco, municipio de Tehuacán, vegetación de matorral, 27-VI-1987, *E. Martínez S. 21690* (ENCB,MEXU).

Distribución y habitat: sólo se conoce de la zona de Tehuacán, en sitios con matorral xerófilo, que prosperan sobre suelos calizos.

Periodo de máxima floración: junio-julio. Periodo de máxima fructificación: septiembre-noviembre.

Karwinskia tehuacana se asemeja a *K. humboldtiana* Zucc., especie ampliamente distribuida en México (Fernández 1988), pero difiere conspicuamente en sus hojas y frutos. Las hojas presentan un color verde limón muy llamativo aún en los ejemplares ya herborizados, las nervaduras secundarias del envés se presentan muy resaltadas y el tamaño de los frutos es notablemente más grande. Es muy probable que estas especies presenten una evolución paralela; sin embargo, en estos momentos es difícil tener un esquema de la filogenia del género *Karwinskia*, esperamos en un futuro cercano tener los elementos necesarios para proponer dicho esquema.

AGRADECIMIENTOS

Se agradece a las autoridades de los herbarios ENCB y MEXU las facilidades prestadas para la consulta del material.

La Dra. Concepción Rodríguez y la Biól. Ma. de la Luz Arreguín tuvieron la amabilidad de leer críticamente el manuscrito.

El biólogo Alfonso Barbosa es el autor de la ilustración de este artículo.

LITERATURA CITADA

- Fernández, N.R. 1988. El Género *Karwinskia* Zucc. (Rhamnaceae) en México. Tesis. M. en C. Esc. Nac. de Ciencias Biológicas. México, D.F. 89 pp.

**OBSERVATIONS ON TOOTHACHE GRASS (*CTENIUM AROMATICUM*
[POACEAE: CHLORIDEAE]) WITH PARTICULAR REFERENCE TO FIRE**

M.H. MacRoberts & B.R. MacRoberts

Bog Research, 740 Columbia, Shreveport, Louisiana 71104 U.S.A.

ABSTRACT

Ctenium aromaticum is fire dependent. In Louisiana we have observed that flowering is confined to the first postfire growing season. Experiments with mechanical removal of litter produced interesting but not consistent results.

KEY WORDS: *Ctenium aromaticum*, bog, fire, Kisatchie National Forest, Poaceae

During our work on Louisiana bogs (MacRoberts & MacRoberts 1988, 1990a, 1991, 1992), we have had the opportunity to observe the effect of fire, usually by way of prescribed dormant season (winter) burns, on various plant species (MacRoberts & MacRoberts 1990b). One of these is a perennial grass, *Ctenium aromaticum* (Walt.) Wood, colloquially known as toothache grass.

Toothache grass is a common bog species. Its range extends along the southern coastal plain from Virginia to Florida, westward to western Louisiana. *Ctenium* is the dominant ground cover in some of the bogs we have studied (MacRoberts & MacRoberts 1991); while in others it is entirely absent (MacRoberts & MacRoberts 1988, 1990a, 1992). It is a conspicuous grass, growing in clumps. The erect flowering culms are up to 140 cm tall with as many as 40 inflorescences per square meter. In some bogs in western Louisiana there are hundreds of thousands of inflorescences.

We first became aware that there was some relationship between fire and *Ctenium* flowering when we spent a couple of days (15 and 16 June 1990) in the pine savannahs near Lake Ramsay, a few miles northwest of Covington, Louisiana. We noted that in a large savannah bisected by a road, flowering *Ctenium* blanketed the side that had been burned the previous winter; whereas on the other side of the road, which had not been burned, *Ctenium* had not put up any new flowering stems. In another instance, on the unburned

side of a fire line through a savannah there were only old *Ctenium* flowering stems; whereas on the other side, which had been burned only a few months previously, *Ctenium* flowered profusely.

In 1991 we were presented with a test of our earlier observations. In a section of the Kisatchie Ranger District of the Kisatchie National Forest, where we do much of our bog research and where all of the bogs have *Ctenium*, the Forest Service regularly sets prescribed burns. In February 1990, they burned three compartments containing twenty-seven bogs. *Ctenium* bloomed profusely in all of these bogs that summer. In early 1991, the Forest Service burned only one of these three compartments containing $14\frac{1}{2}$ bogs (one bog straddles two compartments and had a fire line through it; another bog only partly burned presumably because of high moisture content). Between July 13 and August 3, 1991, we surveyed these twenty-seven bogs.

The results of the survey were clear: *Ctenium* flowered in profusion in all bogs or parts of bogs burned in 1991; only old stems were present in unburned bogs. The bog that straddles two compartments and the bog that only partly burned provide the test. In both cases, *Ctenium* did not flower in the unburned portions but flowered in the burned portions. Counting the half bogs, 14 ($12 + \frac{1}{2} + \frac{1}{2}$) bogs were burned and flowered, while 13 ($12 + \frac{1}{2} + \frac{1}{2}$) were not burned and did not flower.

We have subsequently made many similar observations on toothache grass in the Kisatchie Ranger District and in the Vernon Ranger District. It is such a consistent indicator of fire history that we now use the condition of *Ctenium* flower stalks (since they do not readily drop) as the best indicator for determining when a bog last burned. New flower stalks signify that it was burned since the last growing season, old but standing flower stalks indicate one year past, bedraggled stalks indicate that an area was burned two years ago, and no stalks but leaf clumps that it was burned three or more years ago.

In order to further test the relationship between fire and *Ctenium* (flowering on January 4 and 11, 1992) we established 22 permanent one meter square plots in *Ctenium*-rich areas of eight bogs in the Kisatchie Ranger District. Each plot was sickled and carefully raked to remove litter. In bogs where it is present, *Ctenium* is the major source of litter, which is often so deep that it obscures the ground. All plots were established in bogs that had the same fire history; they had been burned two years ago. Four of the bogs (11 plots) were burned in early February 1992, while the other four bogs (11 plots) were not burned. The nonplot portions of all bogs acted as the control. Our purpose was to see whether the removal of litter without fire would have the same effect as its removal by fire (see Facelli & Pickett 1991).

On June 6, 1992, when the *Ctenium* in the burned bogs was in full flower we examined all of the plots. The *Ctenium* in unburned plots did not put up inflorescences (nor did any of the *Ctenium* elsewhere in these bogs), while *Ctenium* in plots in burned bogs flowered (as did *Ctenium* throughout these

bogs). On June 6, the plots in both burnt and unburnt bogs were open: bare soil and small forbs, such as *Drosera*, *Chaptalia*, *Utricularia*, and *Xyris*, were visible. The absence of litter, therefore, does not seem to be the controlling factor in *Ctenium* flowering. What was somewhat surprising was that the *Ctenium* in the cleared plots in burned bogs flowered even though these plants could not have burned much since most of the litter had been removed. How hot the fire became in such plots is not known, but perhaps close proximity to high temperatures produced by adjacent litter may have helped to produce the effect. Larger plots might have shown a central/peripheral effect (eliminating the possibility of an edge effect) and should be considered in future studies. However, in our experience, one meter plots should be sufficiently large since, in bogs that have not completely burnt, at the burn edge there is always a sharp line of flowering and nonflowering *Ctenium* within inches of one another.

While it is a common observation that winter or spring burns usually stimulate greater growth and flowering in grasses, flowering entirely confined to the first postfire growing season is apparently rare (Biswell & Lemon 1943, Daubenmire 1968, Komarek 1974, Vogl 1974, Knapp & Seastedt 1986, Stolzenburg 1991, Christensen 1988, Robbins & Myers 1989). This apparent dependence in *Ctenium*, insofar as we have been able to discover, has not been previously reported (but see comments by Grelen & Hughes 1984:46).

The relationship of *Ctenium* and fire may not be so simple as this, however. Steve Orzell and Edwin Bridges (pers. comm.) report that, while both nongrowing season and growing season burns produce a dramatic display of flowering in *Ctenium*, they have seen individual plants flower in areas where there has not been any recent fire. While Orzell & Bridges do not give quantitative data, out of the tens of thousands of plants we have observed, we have only seen a few that had bloomed in the absence of fire.

However, let us make one caveat to the above. When we initially established our meter plots to determine if mechanical clearing affected flowering, we established some backup plots on the Vernon Ranger District of the Kisatchie National Forest 60 km to the south of our study plots. We did not re-examine these since they were not needed. But as fate would have it, we had a potted *Ctenium* plant at our home and it produced a single flower stalk in late July. This plant had not been burned. We therefore decided that we should check our plots to see if flowering might be delayed. We examined five of the Vernon Ranger District backup plots on August 4. The bogs in which they were established had not been burned the previous year and there were no new flower stems. But in the five plots, there were 0, 1, 3, 9, and 25 stems in flower. None of these plots showed any ground disturbance additional to the mechanical clearing we had done or any other factors that might account for flowering. Although this is a drastic reduction in flowering as compared to burned plots, it is flowering nonetheless. The next day we re-examined all 11 plots on the Kisatchie Ranger District. None had flowers!

Clearly, additional observations are required to establish the factors involved in stimulating flowering in *Ctenium*. Perhaps there are regional or site specific differences affecting or complicating *Ctenium* flowering as appears to be the case with other grasses (e.g., Robbins & Myers 1989), factors that would be interesting to explore further.

It has long been thought that southeastern bogs and savannahs, like many other plant communities within the longleaf pine ecosystem, are maintained by fire (Folkerts 1982, Smith 1991, Frost *et al.* 1986, Martin & Smith 1991, Platt *et al.* 1988, Robbins & Myers 1989, Bridges & Orzell 1989, Noss 1988, Olsen 1992). The existence of a fire dependent species like *Ctenium aromaticum*, which is fidel to bogs and wet savannahs, certainly strengthens this view.

ACKNOWLEDGMENTS

The continuing cooperation and assistance of the staff of the Kisatchie National Forest have been instrumental in making this study, as in all our botanical work, possible. Latimore Smith and Nelwyn McInnis, Louisiana Natural Heritage Program, directed us to the Lake Ramsay savannahs. D.T. MacRoberts and Jay Wipff commented on an earlier version of this paper. Jay Wipff, Steve Orzell, and Edwin Bridges shared their observations with us.

LITERATURE CITED

- Biswell, H.H. & P.C. Lemon. 1943. Effects of fire upon seedstalk production in range grasses. *J. Forestry* 41:844.
- Bridges, E.L. & S.L. Orzell. 1989. Longleaf pine communities of the west gulf coastal plain. *Natural Areas Journal* 9:246-263.
- Christensen, N.L. 1988. Vegetation of the southeastern coastal plain. In M.G. Barbour & W.D. Billings, eds., *North American Terrestrial Vegetation*. Pp. 318-363. Cambridge University Press, New York, New York.
- Daubenmire, R. 1968. Ecology of fire in grasslands. *Advances in Ecological Research* 5:209-226.
- Facelli, J.M. & S.T.A. Pickett. 1991. Plant litter: Its dynamics and effects on plant community structure. *Botanical Review* 57:1-32.
- Folkerts, G.W. 1982. The gulf coast pitcher plant bogs. *American Scientist* 70:260-267.

- Frost, C.C., J. Walker, & R.K. Peet. 1986. Fire dependent savannas and prairies of the southeast. In D.L. Kulhavy & R.N. Conner, eds., *Wilderness and Natural Areas of the Eastern United States*. P. 348-357. Center for Applied Studies, Stephen F. Austin State University, Nacogdoches, Texas.
- Grelen, H.E. & R.H. Hughes. 1984. Common herbaceous plants of southern forest range. U.S. Dept. of Agriculture, Research Paper SO-210.
- Knapp, A.K. & T.R. Seastedt. 1986. Detritus accumulation limits productivity of tallgrass prairie. *BioScience* 36:662-668.
- Komarek, E.V. 1974. Effects of fire on temperate forests and related ecosystems: southeastern United States. In T.T. Kozlowski & C.E. Ahlgren, eds., *Fire and Ecosystems*. Pp. 251-277. Academic Press, New York, New York.
- MacRoberts, B.R. & M.H. MacRoberts. 1988. Floristic composition of two west Louisiana pitcher plant bogs. *Phytologia* 65:184-190.
- . 1990a. Vascular flora of two west Louisiana pitcher plant bogs. *Phytologia* 68:271-275.
- . 1991. Floristics of three bogs in western Louisiana. *Phytologia* 70:135-141.
- . 1992. Floristics of four small bogs in western Louisiana with observations on species/area relationships. *Phytologia* 73:49-56.
- MacRoberts, M.H. & B.R. MacRoberts. 1990b. Notes on the occurrence of *Platanthera integra* (Nutt.) ex Beck (Orchidaceae) in west central Louisiana. *Phytologia* 69:378-381.
- Martin, D.L. & L.M. Smith. 1991. A survey and description of the natural plant communities of the Kisatchie National Forest Winn and Kisatchie Districts. Unpublished report, Louisiana Natural Heritage Program, Department of Wildlife and Fisheries, Baton Rouge, Louisiana.
- Noss, R.F. 1988. The longleaf pine landscape of the southeast: Almost gone and almost forgotten. *Endangered Species Update* 5(5):1-8.
- Olsen, M.S. 1992. Effects of early and late growing season fires on resprouting of shrubs in Longleaf pine savannas and embedded seepage savannas. M.S. Thesis, Louisiana State University, Baton Rouge, Louisiana.
- Platt, W.J., G.W. Evans, & S.L. Rathbun. 1988. The population dynamics of a long-lived conifer (*Pinus palustris*). *Amer. Nat.* 31:491-525.

- Robbins, L.E. & R.L. Myers. 1989. Seasonal effects of prescribed burning in Florida: A review. The Nature Conservancy. Tallahassee, Florida.
- Smith, L.M. 1991. Louisiana longleaf an endangered legacy. Louisiana Conservationist. May/June 24-27.
- Stolzenburg, W. 1991. The wiregrass mystery. Nature Conservancy 41:28-29.
- Vogl, R.J. 1974. Effects of fire on grasslands. In T.T. Kozowski & C.E. Ahlgren, eds., *Fire and Ecosystems*. Pp. 139-194. Academic Press, New York, New York.

STUDIES ON THE GENUS *BIDENS* L. (COMPOSITAE) FROM THE EASTERN HEMISPHERE. 4. A NEW SPECIES FROM ETHIOPIA

T.G.J. Rayner

Department of Botany, Plant Science Laboratories, University of Reading,
Whiteknights, P.O. Box 221, Reading, RG6 2AS, U.K.

ABSTRACT

A new species, *Bidens kefensis* T.G.J. Rayner, is described from the Kefa administrative region of Ethiopia. It is shown to be a very distinctive taxon, perhaps most closely related to *B. ugandensis* (S. Moore) Sherff from which it is primarily distinguished by its fewer ray florets, annular thickening at the base of the limb of the disc floret corollas, and dorsally produced cartilaginous corona of the cypselas.

KEY WORDS: *Bidens*, Compositae, taxonomy, Ethiopia

During continuing revisional studies of Eastern hemisphere *Bidens*, a collection from Ethiopia with distinctive capitular characters was discovered among specimens housed at MO and WAG. An examination of the revision of *Bidens* in northeastern tropical Africa by Mesfin (1984) has failed to reveal any species possessing the same characters. Thus it is here described as a new species.

Bidens kefensis T.G.J. Rayner, *sp. nov.* TYPE: ETHIOPIA. Kefa administrative region, about 10 km SW of Jima, 18 Dec. 1965, *W.J.J.O. de Wilde, cum suis*, 9266 (HOLOTYPE: WAG; Isotypes: MO, WAG).

Species nova haec similis *Bidenti ugandensi* (S. Moore) Sherff sed ab ea differt plantis tantum usque ad centum viginti centimetra altis, caulibus simplicibus aut ad basem ramosis, foliis omnibus oppositis, laminis foliorum indivisis aut lobatis non pinnatisectis, involucris glabris aut ad basem sparsim hispidis-pubentibus, phyllariis exterioribus uniseriatis linearibus supra parum dilatatis aut linearibus-oblongatis, flosculis radii sex aut septem, corollis flosculorum disci ad basem limbi annulare parum incrassatis,

filis staminum tantum 0.7-1.4 millimetris longis, ramis stigmatum solummodo 1.0-1.2 millimetris longis, cypselis tantum 3.5-4.4 millimetris longis, apice glabris et corona straminea cartilaginea dorsaliter producta usque ad 0.35 millimetri longa 0.7 millimetri lata instructis, aristis cypselarum (ubi adsunt) nudis tantum usque ad 0.4 millimetri longis, carpopodio cartilagineo parum dorsaliter producto usque ad 0.2 millimetri longo basi instructis.

Perennial herbs, to 0.8-1.2 m tall; stems arising from a woody rootstock, sometimes prostrate below, erect above, simple or branched at base; stems and branches more or less terete or subtetragonal above, 1.6-2.4 mm diam. near base, to 0.6-1.3 mm diam. beneath peduncles, striate-sulcate above, somewhat irregularly sulcate below, dark brown toward base, becoming pale green or pale green-yellow above, woody from base up to ca. 1/3 of length of stem, glabrous or, especially above, with isolated, minute (ca. 0.1-0.2 mm long), erect, uniseriate, few-cellular hairs. Leaves decussate, sessile, 0.4-7.4 cm long \times 0.8-6.8 mm wide, the largest generally toward the middle, more or less gradually reduced above and below, linear or narrowly ovate linear to narrowly elliptic in outline; apex acute to subacute, callose indurated; bases narrowly cuneate from about middle of lamina or below, connate; margin entire or serrate to serrate incised or more rarely deeply linear lobed, sometimes slightly revolute or involute and callose indurated; teeth or lobes 1-3 on each side, 0.8-6.2 mm long, acute, callose indurated and slightly mucronate; lamina somewhat coriaceous, pale olive-green, faces glabrous or sparsely hispidulous, margin with more or less 2 rows of antrorse, hispidulous hairs especially in the upper half of the leaf. Capitula radiate, heterogamous, erect, 2.7-3.8 cm diam. \times 6-7 mm high at anthesis, to 9 mm high in fruit, solitary or 2-3 in lax cymes at apices of stems and branches; receptacles flat or slightly convex; peduncles to 2.6-9.8 cm long, very slender, 0.3-0.6 mm diam. at anthesis, to 0.9 mm diam. in fruit, tetragonal to more or less terete, striate-sulcate, glabrous or sparsely pubescent, sometimes somewhat densely pubescent in upper part; ebracteate or with 1-3 alternate bracts resembling the outer phyllaries. Involucre depressed hemispheric, glabrous or sparsely hispid-pubescent at base; outer phyllaries uniseriate, 7-9, linear and slightly dilated above to narrowly obovate linear, with a subacute to obtuse, callose indurated and mucronulate apex, entire at margin, 3.2-3.7 mm long \times 0.7-0.8 mm wide at anthesis, to 4.3 mm long in fruit, erect, green, sometimes darkened at apex, with 1-3 red nerves, glabrous; inner phyllaries uniseriate, 6-7, ovate to ovate elliptic and often attenuate from about middle, subabruptly narrowed near the more or less acute to subobtuse apex, 4.1-5.0 mm long \times 1.6-1.9 mm wide at anthesis, to 6.1 mm long \times to 2.3 mm wide in fruit, erect, membranous, dark brown to stramineous or yellow, darkest toward base and at apex, with 10-15 red-brown nerves, glabrous or dorsal surface sparsely pubescent chiefly along the median

nerve or somewhat densely so especially toward the puberulous apex. Ray florets 6-7, neuter; ovary more or less oblong, 1.0-1.3 mm long \times 0.7-0.8 mm wide, glabrous, exaristate, style absent; corolla tube 1.4-1.8 mm long, glabrous or more or less pubescent; ray yellow or yellow orange, oblong to elliptic or ovate oblong, 1.4-1.5 cm long \times 5.3-5.8 mm wide, with 8-13 red-brown nerves, glabrous; apex rounded or subobtusate, entire or subentire to slightly cuspidate or emarginate, sinus to 0.3 mm deep. Paleae narrowly oblong or narrowly elliptic-oblong, gradually attenuate above to the subacute to obtuse or somewhat rounded and often slightly mucronate apex, entire at margin, 4.5-6.0 mm long \times 0.8-0.9 mm wide at anthesis, to 6.6 mm long in fruit, membranous, glabrous, pale yellow, slightly darkened toward apex, with 1-3 pairs of red or orange brown nerves, especially the central pair darker above, the lateral pairs sometimes coloured only toward middle. Disc florets 22-31; corolla yellow or orange yellow, glabrous or lobes dorsally sparsely pubescent; limb campanulate, 2.4-2.6 mm long \times 1.0-1.4 mm wide, slightly annularly thickened at base, apex (4-)5 lobed; lobes triangular, acute at apex, 0.5-0.6 mm long \times 0.5-0.6 mm wide at base, papillate on margin; limb gradually or subabruptly attenuate below into a narrow, 1.1-1.2 mm long \times 0.4-0.5 mm wide, terete tube; anthers 1.8-2.0 mm long \times 0.5-0.8 mm wide, brown; endothecial tissue with polarized thickening; apical appendages narrowly ovate-triangular, obtuse at apex, 0.4-0.5 mm long \times 0.25-0.30 mm wide, often with a dark brown longitudinal median nerve, margins recurved; basal appendages sagittate, not or just reaching base of the filament collar; collar 0.5-0.6 mm long \times 0.15-0.20 mm wide; filament 0.7-1.4 mm long, involute; style 3.7-5.5 mm long, bulbous or at least slightly dilated at base, with caudate, 1.0-1.2 mm long branches; stylopodium cylindric. Cypselas unwinged but margin usually with a narrow (to 0.05 mm wide) and often interrupted, light brown, cartilaginous extension; body oblong or oblong elliptic, 3.5-4.4 mm long \times 1.3-1.7 mm wide, grey-black, strongly compressed, dorsal face convex or somewhat carinate, ventral face concave with a raised median rib, both faces 4 to 10 striate-sulcate and glabrous or subsparingly erect setose chiefly above, the setae often arising from light brown tubercles, margins more or less densely erect setose, the setae generally becoming slightly longer toward the apex; apex surmounted by a corona, glabrous; corona stramineous, cartilaginous, dorsally produced, slightly flattened, laterally biaristate or rarely exaristate, to 0.10-0.35 mm tall \times 0.5-0.7 mm wide; aristae erect or slightly divergent, rounded-trigonous at base, subulate above, pale yellow, to 0.15-0.40 mm long \times ca. 0.06 mm wide at base, nude; base of cypselas with a short (to 0.1-0.2 mm long), slightly dorsally produced, cartilaginous flaplike carpodium.

Bidens kefensis is apparently endemic to the highlands of southwestern Ethiopia, being known at present only from the type locality near Jima. It was described by de Wilde as growing in a marshy place at an altitude of ca. 1800 m.

It is a distinctive species with no known near relatives in Ethiopia, and is perhaps most closely related to *Bidens ugandensis* (S. Moore) Sherff from western central Africa. *Bidens kefensis*, however, differs from this species in the following ways: stems only to 1.2 m tall (to 2.3 m tall in *B. ugandensis*), simple or branched at base (not sometimes branched above); leaves all opposite (not uppermost sometimes alternate), with lamina undivided or lobed (not often pinnatisect); involucre glabrous or sparsely hispid pubescent at base (not densely hispid at base); outer phyllaries 1 seriate (not 1 to 2 seriate), linear and slightly dilated above or linear-oblongate (not linear to narrowly ovate or narrowly elliptic); ray florets 6-7 (not 8-16); corolla of disc florets slightly annularly thickened at base of limb (not unthickened); filaments of the stamens 0.7-1.4 mm long (not 3.6-4.4 mm long); stylar branches 1.0-1.2 mm long (not 1.4-2.2 mm long); cypselas 3.5-4.4 mm long (not 4.3-14.1 mm long), apex glabrous, with a stramineous, cartilaginous, dorsally produced, to 0.35 mm long \times to 0.7 mm wide corona (not subsparingly to somewhat densely erect setose); aristae to 0.4 mm long, nude (not to 3.1 mm long and often barbed); base of cypselas with a cartilaginous, slightly dorsally produced, to 0.2 mm long carpopodium (not hollow and encircled by a cartilaginous, to 0.4 mm long carpopodium).

ACKNOWLEDGMENT

I am most grateful to C. Jeffrey (K) for correcting the Latin diagnosis.

BIBLIOGRAPHY

- Mesfin Tadesse. 1984. The genus *Bidens* (Compositae) in NE tropical Africa. Symb. Bot. Upsal. 24(1):138 pp.

THE GOLDENASTERS OF CALIFORNIA, *HETEROTHECA* (COMPOSITAE: ASTEREA): NEW NAMES AND COMBINATIONS

John C. Semple

Department of Biology, University of Waterloo, Waterloo, Ontario,
CANADA N2L 3G1

ABSTRACT

Extensive field work and herbarium specimen studies on the goldenasters of California indicated that a number of new names and combinations are required. The following new names and combinations are proposed: *Heterotheca sessiliflora* (Nutt.) Shinnery ssp. *bolanderi* (A. Gray) Semple, *H. sessiliflora* ssp. *echioides* (Benth.) Semple, *H. sessiliflora* ssp. *echioides* var. *bolanderioides* Semple, *H. sessiliflora* ssp. *echioides* var. *camphorata* (Eastwood) Semple, *H. sessiliflora* ssp. *fastigiata* (E. Greene) Semple, *H. sessiliflora* ssp. *fastigiata* var. *sanjacintensis* Semple, *H. villosa* (Pursh) Shinnery var. *scabra* (Eastwood) Semple, *H. villosa* var. *shevockii* Semple.

KEY WORDS: *Heterotheca*, Compositae, Astereae, California, Goldenasters

The following new combinations and names are required for use in the forthcoming *The Jepson Manual, Higher Plants of California* (J.C. Hickman, ed. 1993). Justification for the nomenclatural changes and additions will be presented in full elsewhere. Nomenclatural changes are based on examination of hundreds of herbarium specimens, field work conducted over a ten year period, and multivariate morphometric analyses of the *Heterotheca sessiliflora* and *H. villosa* complexes. All specimens cited below, including types, were examined during the course of the study.

Heterotheca sessiliflora (Nutt.) Shinnery ssp. *bolanderi* (A. Gray) Semple, *comb. et stat. nov.* BASIONYM: *Chrysopsis bolanderi* A. Gray, Proc. Amer. Acad. Arts 6:543. 1866. *Chrysopsis villosa* (Pursh) Nutt.

var. *bolanderi* (A. Gray) A. Gray, *Syn. Fl. N. Amer.* 1(2):123. 1884. *Heterotheca bolanderi* (A. Gray) Harms, *Brittonia* 26:61. 1974. TYPE: U.S.A. California: Oakland Hills near San Francisco, 1863, *Bolander* 2466 (HOLOTYPE: GH; Isotypes: K,US).

***Heterotheca sessiliflora* (Nutt.) Shinnery ssp. *echioides* (Benth.) Semple, *comb. et stat. nov.* BASIONYM: *Chrysopsis echioides* Benth., *Bot. Voy. Sulphur* 25. 1844. *Chrysopsis villosa* (Pursh) Nutt. var. *echioides* (Benth.) A. Gray, *Syn. Fl. N. Amer.* 1(2):123. 1884. *Heterotheca echioides* (Benth.) Shinnery, *Field & Lab.* 19:71. 1951. TYPE: U.S.A. California: Bodegas, *Hinds s.n.* (HOLOTYPE: K).**

***Heterotheca sessiliflora* (Nutt.) Shinnery ssp. *echioides* (Benth.) Semple var. *bolanderioides* Semple, *var. nov.* TYPE: U.S.A. California: Contra Costa Co., Charles Tilden Reg. Park, Vollmer Peak. Scattered population along ridge top, in rocks and loose gravel in grassy area between shrubs, 16 Aug 1990, 2n = 36, *Semple, Suripto, & Ahmed* 9399 (HOLOTYPE: UC; Isotypes: CAN,CAS,MO,NY,RSA,WAT).**

Heterotheca sessiliflora (Nutt.) Shinnery ssp. *echioides* (Benth.)
Semple var. *echioides* *accedens* sed foliis intermediisque interdum dense strigosibus longibus, involucri grandibus, capitulescentiis corymbiformibus compactibus.

Perennial from stout woody taproots, the stems several to many, ascending-erect, 17-45 cm tall, sparsely to densely strigose-hispid, the hispid hairs fewer than the shorter appressed ones, sometimes becoming more glandular and sparsely pubescent above. Lower stem leaves oblanceolate, 15-45 mm long, 4.5-10. mm wide, subpetiolate to sessile, cuneate, acute, mucronate, moderately to densely hispid-strigose on both surfaces; margins entire, strigose, longest hispid hairs near base, not undulate. Upper stem leaves lanceolate to elliptic, sessile, slightly reduced upward, 11-35 mm long, 3.5-7.5 mm wide, sparsely to densely villous-strigose (5-80 hairs/mm²), moderately to densely glandular (7-40 glands/mm²), acute to obtuse, mucronate. Capitulescence cymose-paniculiform, heads (1-)4-16; peduncles as upper stems to more glandular, bracts few, lower ones oblanceolate, like leaves, sometimes reduced upward to < 5 mm, those immediately below head 4-11 mm long, 0.8-4.0 mm wide. Involucres cylindrical to campanulate when fresh, campanulate-hemispheric upon drying, (7-)8-11(-12) mm tall; phyllaries in 4-5 imbricate series, outer 1/4-1/3 length of inner, narrowly triangular; mid series lanceolate, sparsely

to moderately glandular, sparsely to moderately strigose, margins hyaline, fimbriate-ciliate apically; inner ones similar. Ray florets 7-16, strap yellow, 4.5-10. mm long, 0.7-2.4 mm wide. Disc florets 28-70, yellow, glabrous, corolla barely ampliate, 5.5-7.5 mm long, lobes 0.4-1.0 mm long, sparsely strigose, hairs 0.3-0.7 mm long. Achenes 3-4 mm long, moderately to densely strigose; pappus off white, double, outer whorl of a few linear scales 0.25-0.50 mm long, inner whorl of 35-45 barbellate bristles 5.5-8.0 mm long. Chromosome number: $2n = 36$.

PARATYPES: U.S.A. California: Santa Clara Co.: Page Mill Rd. NE of CA-35, 21 Sep 1987, *Semple & Chmielewski 8915* (WAT). Marin Co.: San Geronimo, grass covered rocky ridge N. of golf course, common on slope above and below Nicasio valley road, large number of plants, 15 Aug 1990, *Semple, Suropto, & Ahmed 9893* (WAT). Santa Cruz Co.: W of Palo Alto, CA-35 17.5 km SE of CA-84, *Semple & Chmielewski 8918* (WAT); CA-35 NW of Saratoga, high elevation, *Semple & Semple 5670* (JCS-personal herbarium, MO, MT, UC, USF, WAT). [Additional duplicates to be distributed].

Variety *bolanderioides* is endemic to serpentine soils mostly on hill and mountain tops surrounding San Francisco Bay. As the name indicates, it is similar to *ssp. bolanderi* in which most collections of the taxon have been placed in the past. Individuals of *var. bolanderioides* can be similar in indument to diploid (rarely tetraploid) *var. camphorata*, which has relatively few hairs (for the *ssp.*) on its upper stem and rameal leaves, and to diploid *var. echioides*, which usually has a leaf indument of dense hispid and strigose hairs obscuring the underlying glands and smaller involucres. It is uncertain whether *var. bolanderioides* is a tetraploid derivative of *var. echioides* or whether it may be an allopolyploid involving *ssp. echioides* and *ssp. bolanderi*.

Heterotheca sessiliflora (Nutt.) Shinnars *ssp. echioides* (Benth.) Semple *var. camphorata* (Eastwood) Semple, *var. nov.* BASIONYM: *Chrysopsis camphorata* Eastwood, Zöe 5:81. 1900. *Chrysopsis villosa* (Pursh) Nutt. *var. camphorata* (Eastwood) Jepson, *Man. Fl. Pl. Calif.* 1036. 1925. *Heterotheca camphorata* (Eastwood) Semple, *Canad. J. Bot.* 58:148. 1980. TYPE: U.S.A. California: Santa Cruz Co.; Glenwood, Jul 1900, *Davis s.n.* (HOLOTYPE: CAS; Isotypes: DS, GH(2), NY(2), RM(3), UC, US).

Heterotheca sessiliflora (Nutt.) Shinnars *ssp. fastigiata* (E. Greene) Semple, *comb. et stat. nov.* BASIONYM: *Chrysopsis fastigiata* E. Greene, *Pittonia* 3:296. 1898. *Chrysopsis villosa* (Pursh) Nutt. *var. fastigiata* (E. Greene) H.M. Hall, *Univ. Calif. Publ. Bot.* 3:43. 1907. *Heterotheca fastigiata* (E. Greene) Harms, *Brittonia* 26:61. 1974. TYPE:

U.S.A. California: San Bernardino Mts, 1000-1500' [not 10000-15000 as in protologue], 15 Oct 1895, *Parish 3815* (HOLOTYPE: NDG; Isotypes: CAS,GH,UC,US).

Heterotheca sessiliflora (Nutt.) Shinnery ssp. *fastigiata* (E. Greene) Semple var. *sanjacintensis* Semple, var. nov. TYPE: U.S.A. California: Riverside Co., CA-243 just S of Idyllwild Park at Manzanita Drive, 30 Sep 1987, $2n = 9_{II}$, *Semple & Chmielewski 8982* (HOLOTYPE: WAT; Isotypes (all shoots from same plant): CAS,MT,NY,RSA,UC).

Heterotheca sessiliflora (Nutt.) Shinnery ssp. *fastigiata* (E. Greene) Semple var. *fastigiata* accedens sed foliis utrinque sparsusque vel intermediusque interdum densis strigosibus et hispidibus, utrinque intermediusque vel densis glandulibus, viridibus non albis nec canescentibus.

Perennial from stout woody taproots, the stems several to many, ascending-erect, 35-105 cm tall, moderately appressed strigose-hispid (hairs often broken off), becoming densely glandular and moderately pubescent above. Lower stem leaves oblanceolate, 12-38 mm long, 3-8 mm wide, subsessile, cuneate, acute, moderately (rarely densely) hispid-strigose on both surfaces; margins entire, strigose, longer hispid hairs near base, undulate. Upper stem leaves lanceolate, sessile, reduced upward, 11-20 mm long, 3.5-6.5 mm wide, sparsely to densely glandular (6-42 glands/mm²), sparsely to densely short strigose (27-150 hairs/mm²), margins usually distinctly undulate. Capitulescence cymose-paniculiform, branches ascending, heads 5-60. Peduncles densely glandular, bracts few, lower ones lanceolate, leaflike, reduced upward to 0.5-1.5 mm long, 0.3-1.0 mm wide, phyllarylike. Involucres cylindrical to turbinate when fresh, campanulate upon drying, 7.5-12. mm high; phyllaries in 5-6 imbricate series, outer 1/5-1/4 length of inner, mid series narrowly triangular, moderately glandular, very sparsely strigose, margins hyaline, fimbriate-ciliate apically. Ray florets 5-13, strap yellow, 3.5-5.0 mm long, 0.8-1.8 mm wide. Disc florets 25-45, yellow, corolla somewhat ampliate, 5.3-7.5 mm long, lobes 0.4-0.8 mm long, sparsely pilose, hairs 0.25-0.50 mm long. Achenes 2-3 mm long, moderately strigose; pappus off white, double, outer whorl of a few linear scales 0.25-0.50 mm long, inner whorl of 25-40 barbellate bristles 6-8 mm long. Chromosome number: $2n = 18$.

PARATYPES: U.S.A. California: Riverside Co.: CA-243 just S of Idyllwild Park at Manzanita Drive, 30 Sep 1987, $2n = 9_{II}$, *Semple & Chmielewski 8981* (Each shoot from a separate plant: CAN,CAS,DAO,JCS-personal herbarium,MO,OBI,RM,RSA,SD,UC,WAT).

Variety *sanjacintensis* is endemic to the San Jacinto Mountains and Mt. Palomar areas of southern California. It has a capitulescence form and very undulate stem leaves that are similar to var. *fastigiata*, but its leaves are far less densely strigose than those of var. *fastigiata* and the hairs are somewhat longer. Individuals with less undulate leaves are similar to var. *camphorata* of ssp. *echioides*, which occurs much farther north in California in the Coastal Range from San Mateo to San Luis Obispo counties.

Heterotheca villosa (Pursh) Shinnery var. ***scabra*** (Eastwood) Semple, *comb. nov.* BASIONYM: *Chrysopsis villosa* (Pursh) Nutt. var. *scabra* Eastwood, Proc. Calif. Acad. Sci., ser. 2. 6:294. 1896. TYPE: U.S.A. Utah: San Juan Co., near head of Willow Creek, *Eastwood s.n.* (HOLOTYPE: CAS).

Chrysopsis viscida (A. Gray) E. Greene ssp. *cinerascens* S.F. Blake, Proc. Biol. Soc. Wash. 35:173. 1922. *Heterotheca horrida* (Rydb.) Harms ssp. *cinerascens* (S.F. Blake) Semple, Brittonia 39:381. 1987. TYPE: U.S.A. Utah: Beaver Canyon, among rocks in the oak region, 2 Sept 1909, *Tidestrom 2873* (HOLOTYPE: US).

Semple (1987) discussed this taxon under the synonym *Heterotheca horrida* ssp. *cinerascens*. Additional work on the genus and the *villosa* complex (Semple 1990) resulted in treatment of *H. horrida* as a synonym of *H. villosa* var. *hispida* (Hook.) Harms. At the varietal level Alice Eastwood's epithet has priority over a combination based on Blake's epithet.

Heterotheca villosa (Pursh) Shinnery var. ***shevockii*** Semple, *var. nov.* TYPE: U.S.A. California: Kern Co., Kern R. Canyon, Bodfish, CA-178 at Bodfish-Havilah Rd., off of off-ramp, ca. 850 m el., 16 Nov 1981, *Shevock 9110* (HOLOTYPE: CAS; Isotype: WAT).

Heterotheca villosa (Pursh) Shinnery var. *hispida* (Hook.) Harms *accedens* sed *caulis altae*, *foliae deltoideae-lanceolatae*, *glandulissimae*, *acutae*, *marginibus plerumque involutibus*, *involucellis grandibus*; *chromosomatum numerus* $2n = 36$.

Perennial from stout woody taproots, the stems several to many, ascending-erect, 25-135 cm tall, sparsely strigose, moderately hispid (hairs often broken off in older stems), becoming densely glandular and sparsely hispid-strigose above. Lower stem leaves oblanceolate to lanceolate, 25-55 mm long, 8-26 mm wide, subpetiolate to sessile, cuneate, acute, mucronate, moderately hispid-villous on both surfaces; margins entire, strigose, longer hispid hairs near base.

Upper stem leaves linear lanceolate to lanceolate, sessile, base abruptly tapering, reduced upward, densely glandular, sparsely villous-strigose. Rameal leaves much reduced upward, becoming linear to linear oblanceolate. Capitulescence cymose-paniculiform, heads 3-70, branches elongated in robust shoots. Peduncles long, densely glandular, sparsely hispid-strigose, bracts few, lower ones leaflike, greatly reduced upward, becoming linear to oblanceolate, margins with long hispid hairs. Involucres cylindrical to turbinate when fresh, campanulate upon drying, 9-13 mm tall; phyllaries in 5-6 imbricate series, outer 1/5-1/4 length of inner, narrowly triangular, densely glandular, sparsely strigose especially along the pronounced midvein, margins hyaline, fimbriate-ciliate apically; midseries linear lanceolate to linear oblanceolate, moderately hispid apically, margins similar to outer series. Ray florets 9-14(-18), strap yellow, 5-10 mm long, 0.8-2.0 mm wide. Disc florets (31-)41-68(-77), yellow, glabrous, corolla barely ampliate, 5.4-7.5 mm long, lobes (0.4-)0.5-0.9(-1.1) mm long, glabrous or very sparsely strigose, hairs 0.04-0.27 mm long. Achenes 3.5-4.5 mm long, moderately strigose; pappus off white, double, outer whorl of a few linear scales 0.25-0.50 mm long, inner whorl of 35-45 barbellate bristles 5-7 mm long. Chromosome number: $2n = 36$.

PARATYPES: U.S.A. California: Kern Co., Kern R. Canyon, CA-178 8.2 km SE of CA-155, edge of rd. below rock cut, 26 Sept 1987, *Semple & Chmielewski 8953* (CAS, JCS-personal herbarium, NY, RSA, UC, WAT); CA-178 25.4 km SW of CA-155, Democrat Hot Springs, Democrat Raft Removal Area, 26 Sept 1987, *Semple & Chmielewski 8954* (CAS, RM, RSA, WAT); just E of Miracle Hot Springs, Sequoia Nat'l For. - NE end of Hobo Campground, 18 Aug 1990, *Semple, Suropto, & Ahmed 9363* (WAT); CA-178, ca. 1.5 mi E of Rich Bar, 1000 ft. el., 16 Nov 1981, *Shevock 9105* (CAS). [Additional duplicates to be distributed].

Shevock's Goldenaster is named for James Shevock, an expert on the flora of Kern County, California, who collected the taxon several times in the Kern River Canyon area and who provided valuable personal communication on its habit and habitat. The var. *shevockii* is distinguished from other races of the species by its usually tall stems and lanceolate-deltoid leaves with inrolled margins. Smaller plants are similar to *Heterotheca villosa* var. *scabra* to which var. *shevockii* is undoubtedly closely related. In California, var. *scabra* is known only from a few locations in the Little San Bernardino Mountains (1200-1300 m el.), var. *hispida* occurs in the Sierra Nevadas, the Cascade Mountains and on lava flows in Lassen and Modoc counties (600-3100 m el.), while var. *shevockii* is endemic to the Kern River Canyon (400-900 m el.) in the Greenhorn Mountains. As with all races of *H. villosa* (Semple 1990), in typical form var. *shevockii* is readily recognized, but depauperate and atypical individuals are not easily separated from similar taxa, i.e., var. *hispida* and var. *scabra*.

ACKNOWLEDGMENTS

This research was supported by Natural Sciences and Engineering Research Council of Canada Operating Grantings to the author. The assistance in the field of J. Chmielewski, B.A. Suropto, and B. Semple is gratefully acknowledged. Dr. Judith Canne-Hillicker and Dr. Nag Raj are thanked for their help with the manuscript.

LITERATURE CITED

- Hickman, J.C. , ed. 1993. *The Jepson Manual, Higher Plants of California*. University of California Press. in press.
- Semple, J.C. 1987. New names, combinations and lectotypifications in *Heterotheca* (Compositae: Astereae). *Brittonia* 39:379-386.
- . 1990. Neotypification of *Amellus villosus*, the identity of a Bradbury collection, and typification of some other goldenasters (Compositae: Astereae). *Brittonia* 42:221-228.

BOISDUVALIA, A COMA-LESS EPILOBIUM (ONAGRACEAE)

Peter C. Hoch & Peter H. Raven

Missouri Botanical Garden, P.O. Box 299, St. Louis, Missouri 63166 U.S.A.

ABSTRACT

Evidence from independent analyses of morphological and molecular variation in tribe Epilobieae shows no support for the continued recognition of *Boisduvalia* Spach. Absence of seed comas, the only consistent diagnostic feature of the genus relative to *Epilobium*, now appears to be a secondary loss; other characters reveal a close relationship of *Boisduvalia* to taxa within *Epilobium*. Therefore, all species of *Boisduvalia* are transferred to *Epilobium*, in order better to reflect phylogeny.

KEY WORDS: Onagraceae, *Epilobium*, *Boisduvalia*, New World

Tribe Epilobieae (Munz 1965; Raven 1976) is marked within Onagraceae as monophyletic by probable base chromosome number of $x = 9$ (Raven 1976, 1979), commissural stigmas (Eyde 1982), and dry type stigma surface (Heslop-Harrison 1990); most species (excluding mainly *Epilobium* sect. *Chamaenerion*) also have pollen released in tetrads (Skvarla *et al.* 1978). The tribe traditionally has comprised *Epilobium* (including sections *Chamaenerion* and more recently *Zauschneria*), characterized by the synapomorphy of seed comas (tufts of hair on the chalazal end of the seed), and *Boisduvalia* (Raven & Moore 1965), an entirely annual genus that lacks comas. Recent phylogenetic analyses of the tribe using either morphological data (Hoch & Crisci, in prep.) or variation in chloroplast DNA (Baum *et al.*, in prep.) have revealed relationships among the taxa that are not reflected in the current taxonomy. Even though the exact relationships proposed among the taxa differ between these two studies, both demonstrate convincingly that recognition of *Boisduvalia* as a genus is not supported and that continuing to recognize it renders *Epilobium* paraphyletic.

These studies strongly suggest that *Boisduvalia* evolved from taxa with a coma. By analogy in support of that suggestion, two species of *Epilobium* and populations of a third have secondarily lost their comas (Munz 1965; Raven

& Raven 1976; Seavey *et al.* 1976). Other characters that mark species or groups of species in *Boisduvalia* include capsule specializations and seed shape, but these may be associated with the loss of the coma and may be viewed more accurately as apomorphies within the group, rather than plesiomorphies within the tribe. Consequently, we here propose that *Boisduvalia* be placed in synonymy with *Epilobium*, and make the necessary nomenclatural changes, in order to make the names available for floristic treatments in progress.

The following nomenclatural changes include only names relevant to the transfers. For more complete synonymy of *Epilobium* and the sections, see Raven (1976); for complete synonymies of all species being transferred from *Boisduvalia*, see Raven & Moore (1965).

Epilobium L., *Sp. Pl.* 347. 1753.

Epilobium sect. **Boisduvalia** (Spach) Hoch & Raven, *comb. nov.* **BA-SYNONYM:** *Boisduvalia* Spach, *Hist. Nat. Vég.* 4:383. 1835. *Oenothera* L. sect. *Boisduvalia* (Spach) Torr. & A. Gray, *Fl. N. Amer.* 1:505. 1840. *Oenothera* group *Boisduvalia* (Spach) H. Lév., *Monogr. Oenothera* 296. 1908. **TYPE:** *Epilobium concinnum* (D. Don) Hoch & Raven.

1. **Epilobium concinnum** (D. Don) Hoch & Raven, *comb. nov.* **BA-SYNONYM:** *Oenothera concinna* D. Don in Sweet, *Brit. Fl. Gard.* II, pl. 183. 1833. *Boisduvalia concinna* (D. Don) Spach, *Hist. Nat. Vég.* 4:384. 1835. *Oenothera subulata* (Ruiz & Pavón) H. Lév. *race concinna* (D. Don) H. Lév., *Monogr. Oenothera* 298. 1908. **TYPE:** Sweet, *Brit. Fl. Gard.* II, pl. 183. 1833; plant raised from seeds sent from Chile by H. Cuming (LECTOTYPE, designated by Raven & Moore [1965]).

Boisduvalia subulata (Ruiz & Pavón) Raim. in Engl. & Prantl, *Nat. Pflanzenfam.* III. 7:212. 1893. *Oenothera subulata* Ruiz & Pavón, *Fl. Peruv. Prodr.* 3:82, pl. 316. 1802; non *Epilobium subulatum* (Hausskn.) Rydb., 1913.

2. **Epilobium densiflorum** (Lindl.) Hoch & Raven, *comb. nov.* **BA-SYNONYM:** *Oenothera densiflora* Lindl., *Bot. Reg.* 19: pl. 1593. 1833. *Boisduvalia douglasii* Spach, *Hist. Nat. Vég.* 4:385. 1835, pro syn. *Boisduvalia densiflora* (Lindl.) S. Watson, *Bot. California* 1:233. 1876. **TYPE:** Lindl., *Bot. Reg.* 19: pl. 1593. 1833; plant raised from seeds sent from "Northern California" by Douglas in 1831 (LECTOTYPE, designated by Raven & Moore [1965]).

3. **Epilobium pallidum** (Eastwood) Hoch & Raven, *comb. nov.* BASIONYM: *Boisduvalia pallida* Eastwood, *Leaf. W. Bot.* 2:54. 1937. TYPE: U.S.A. California: Modoc Co., Goose Valley, George Dillman Ranch, 11 July 1912, *Eastwood 1021* (HOLOTYPE: CAS 243301; Isotypes: GH,POM,US).

Boisduvalia macrantha A. Heller, *Muhlenbergia* 2:101. 1905; non *Epilobium macranthum* Hook., 1840.

4. **Epilobium torreyi** (S. Watson) Hoch & Raven, *comb. nov.* BASIONYM: *Oenothera torreyi* S. Watson, *Proc. Amer. Acad. Arts* 8:600. 1873. *Boisduvalia torreyi* (S. Watson) S. Watson, *Bot. California* 1:233. 1876. TYPE: U.S.A. California: Santa Clara Co., New Almaden, 1865, *Torrey 109* (LECTOTYPE, designated by Munz (1941): GH; Isolectotype: NY).

Boisduvalia stricta (A. Gray) E. Greene, *Fl. Francisc.* 225. 1891.

Gayophytum strictum A. Gray, *Proc. Amer. Acad. Arts* 7:340. 1868; non *Epilobium strictum* Muhl., 1813.

- Epilobium sect. Currania** (Munz) Hoch & Raven, *comb. nov.* BASIONYM: *Boisduvalia* Spach. sect. *Currania* Munz, *Darwiniana* 5: 127. 1941. TYPE: *Epilobium cleistogamum* (Curran) Hoch & Raven, selected by Raven & Moore (1965:251).

5. **Epilobium cleistogamum** (Curran) Hoch & Raven, *comb. nov.* BASIONYM: *Boisduvalia cleistogama* Curran, *Bull. Calif. Acad. Sci.* 1:12. 1884. *Oenothera cleistogama* (Curran) H. Lév., *Monogr. Oenothera* 304, 312. 1908. TYPE: U.S.A. California: Solano Co., near Elmira, "May" 1883, *Curran s.n.* (LECTOTYPE, designated by Raven & Moore (1965), CAS 126; Possible isolectotype: GH).

6. **Epilobium pygmaeum** (Speg.) Hoch & Raven, *comb. nov.* BASIONYM: *Oenothera pygmaea* Speg., *Anales Soc. Ci. Argent.* 48:46. 1899. *Boisduvalia pygmaea* (Speg.) Munz, *Physis* 11:278. 1933. TYPE: ARGENTINA. Chubut, Chonkenk-aik, 1 Aug. 1897, *Ameghino s.n.* (HOLOTYPE: LPS not seen).

- Boisduvalia glabella* (Nutt.) Walp., Repert. Bot. Syst. 2:89. 1843.
Oenothera glabella Nutt. in Torr. & A. Gray, Fl. N. Amer. 1:505.
1840; non *Epilobium glabellum* G. Forst., 1786.

ACKNOWLEDGMENTS

We thank the National Science Foundation for support of this work through BSR-8906848. We also thank Kenneth Sytsma and Divide Baum for sharing unpublished results of their molecular analyses, Jorge V. Crisci for collaboration on the morphological phylogeny, and Paul Berry for comments on the manuscript.

LITERATURE CITED

- Eyde, R.H. 1982. Evolution and systematics of the Onagraceae: Floral anatomy. Ann. Missouri Bot. Gard. 69:735-747.
- Heslop-Harrison, Y. 1990. Stigma form and surface in relation to self-incompatibility in the Onagraceae. Nordic J. Bot. 10:1-19.
- Munz, P.A. 1941. A revision of the genus *Boisduvalia* (Onagraceae). Darwiniana 5:124-153.
- . 1965. Onagraceae. N. Amer. Fl., Ser. 2, 5:198-231.
- Raven, P.H. 1976 [1977]. Generic and sectional delimitation in Onagraceae, tribe Epilobieae. Ann. Missouri Bot. Gard. 63:326-340.
- . 1979. A survey of reproductive biology in Onagraceae. New Zealand J. Bot. 17:575-593.
- & D.M. Moore. 1965. A revision of *Boisduvalia* (Onagraceae). Brittonia 17:238-254.
- & T.E. Raven. 1976. The genus *Epilobium* in Australasia. New Zealand Dept. Sci. Industr. Res. Bull. 216:1-321.
- Seavey, S.R., R.E. Magill & P.H. Raven. 1976 [1977]. Evolution of seed size, shape and surface architecture in the tribe Epilobieae. Ann. Missouri Bot. Gard. 64:18-47.
- Skvarla, J.J., P.H. Raven, W.F. Chissole, & M. Sharp. 1978. An ultrastructural study of viscin threads in Onagraceae pollen. Pollen & Spores 20:5-123.

A NEW SPECIES OF *EPILOBIUM* (ONAGRACEAE) IN CALIFORNIA

Peter C. Hoch

Missouri Botanical Garden, P.O. Box 299, St. Louis, Missouri 63166 U.S.A.

ABSTRACT

Epilobium howellii, a new species from California, is described, and compared with similar species in the region. An uncommon plant marked by glandular hairs all over its stems, it is restricted to wet, possibly boggy, areas of the Sierra Nevada Mountains.

KEY WORDS: Onagraceae, *Epilobium*, California

A specimen of *Epilobium* collected in 1975 at Yuba Pass in the northern Sierra Nevada Mountains of California, and recollected in 1981 to obtain seeds for experimental cultivation, proved to be, surprisingly, an undescribed species. A search through hundreds of personal collections and thousands of herbarium specimens on loan for ongoing monographic work on North American *Epilobium* turned up a few additional specimens from elsewhere in the Sierra Nevada, which helped to establish the distinctness of the new taxon. A thorough search of the literature established that it has not been described before. Because the morphological distinctions among many species of *Epilobium* are fine, clear taxonomic keys are few, and the genus has a reputation (sometimes overstated) for hybridization, a conservative taxonomic approach to the delimitation of species in the genus (Trelease 1891; Raven & Raven 1976) is clearly warranted. Nevertheless, observations of the plant in the field, the herbarium, and the greenhouse support recognition of the following new species of *Epilobium*, which would be placed into section *Epilobium*:

Epilobium howellii* Hoch, *sp. nov. TYPE: U.S.A. California: Sierra Co., Yuba Pass summit, along CA Highway 49; 39° 39' N, 120° 30' W; scattered population among grasses and moss in semi-open *Salix* swales; elev. 2,040 m; 27 Jul 1975, Hoch 665 (HOLOTYPE: MO).

Herba perennis tenuis, brevibus filiformibus foliosis stolonibus; caulis undique glandulosus; folia sessilia, 4-20 mm longa, rotundata vel lanceolata; petala alba, 2-3 mm; stigma capitatum; capsulae 35-45 mm longae, subglabrae, pedicelli 25-40 mm; semina 0.8-1.1 mm longa, vix papillosa.

Delicate perennial herb, forming short threadlike stolons with scattered minute leaves. Stems 8-20 cm tall, densely glandular, terete, loosely clumped. Leaves sessile, the blades 4-20 mm long, round to lanceolate or narrower above, tip obtuse to subacute above, margins finely denticulate, stigillose mainly on veins or all over on upper leaves. Inflorescence erect. Flowers small, subcleistogamous; floral tube 0.4-0.8 mm deep; sepals 1.5-2.0 mm long; petals 2-3 mm long, white; stamens in two unequal sets, the longer ones shedding pollen onto capitate stigma prior to petal expansion. Capsules 35-45 mm long, subglabrous, on pedicels 25-40 mm long. Seeds 0.8-1.1 mm long, the surface low papillate; coma dingy, easily detached. $2n = 36$.

Distribution: Scattered locations in the high Sierra Nevada Mountains of California, in Fresno, Mono, and Sierra counties; in mossy meadows and swales, at 2000-2700 m elevation.

Phenology: Flowering period: July-early August; fruiting period: August-October.

Representative specimens examined: U.S.A. California: Fresno Co.: 3 mi. E of Huntington Lake, *Hoch 486* (MO, RSA); 1 mi. SW of Portal Forebay, *Hoch 503* (MO). Mono Co.: Twin Lakes, south shore, *Hoch 533* (MO). Sierra Co.: Yuba Pass summit, *Wagner 4550* (MO).

Chromosome count: $n = 18$, count provided by Warren L. Wagner, from plants cultivated at Missouri Botanical Garden, *Hoch M2616* (MO), from seed source: *Wagner 4550*.

With great respect, I name this plant after John Thomas Howell, enthusiastic collector of Californian epilobiums and observant student of the Sierran flora, in honor of his 90th birthday. This species is described now in order to make the name available for the treatment of *Epilobium* in *The Jepson Manual: Higher Plants of California*.

Epilobium howellii has a similar small, delicate stature, with spreading threadlike stolons, to that of *E. oregonense* Hausskn., but differs from it and most other North American species of *Epilobium* (Hoch 1993; Munz 1965) in having stems covered with glandular pubescence. The flowers are strikingly and consistently small (petals not more than 3 mm long), possibly cleistogamous, and the leaves never more narrow in shape than lanceolate. This combination of features, especially the unusual pattern of pubescence, distinguishes this species also from the so called Alpinae group (Haussknecht 1884), which are of similar habit and stature.

ACKNOWLEDGMENTS

I thank the National Science Foundation for support of this work through BSR89-06848, Warren L. Wagner for recollection of the type population in 1981 to obtain seeds, and for the chromosome count, and Peter H. Raven for his continued support, encouragement, and useful discussion of this work.

LITERATURE CITED

- Haussknecht, C. 1884. *Monographie der Gattung Epilobium*. Jena, Germany.
- Hoch, P.C. 1993 (in press). *Epilobium*, in J.C. Hickman *et al.* (eds.), *The Jepson Manual: Higher Plants of California*. University of California Press, Berkeley, California.
- Munz, P.A. 1965. Onagraceae. *N. Amer. Fl.*, Ser. 2. 5:198-231.
- Raven, P.H. & T.E. Raven. 1976. The genus *Epilobium* in Australasia. New Zealand D.S.I.R. Res. Bull. 216:1-321.
- Trelease, W. 1891. A revision of the American species of *Epilobium* occurring north of Mexico. *Ann. Rep. Missouri Bot. Gard.* 2:69-117.

ERRATUM

In formatting a recent article G.L. Nesom for printing (Taxonomic notes on *Erigeron* [Asteraceae: Astereae] of California, Nevada, and Arizona. *Phytologia* 73:186-202. 1992), the editor made changes (p. 198) in the submitted manuscript that did not conform to the intent of the author. Following is the version as originally submitted by Nesom, with the new combination intended to place *E. austiniae* at equivalent (varietal) rank with *E. var. chrysopsidis* and *E. var. brevifolius*, and which makes only the claim that a trinomial is being validated (see ICBN Art. 24). The new varietal combination can still be regarded as validly published in *Phytologia* 73:198. 1992.

"*Erigeron chrysopsidis* A. Gray

Subsp. *chrysopsidis*

Var. *chrysopsidis*

Var. *brevifolius* Piper, Bull. Torrey Bot. Club 27:395. 1900.

TYPE: Oregon, [Wallowa Co.,] subalpine ridge of the Wallowa Mts. near the [Wallowa] Lake, 29 Jul 1899, *W. C. Cusick* 2270 (Holotype: WS; isotypes: GH!, MO!, NY!).

Subsp. *austiniae* (E. Greene) Cronq., Brittonia 6:196. 1947. Based on *E. austiniae* E. Greene (below)

Var. *austiniae* (E. Greene) Nesom, comb. et stat. nov. *Erigeron austiniae* E. Greene, Erythea 3:100. 1895. TYPE: California, Modoc County, Davis Creek, May 1894, *Mrs. R. M. Austin s.n.* (Lectotype, designated here: ND-G; isolectotypes: NY!, PH, UC)."

BOOKS RECEIVED

Agricultural Plants, Second Edition. R.H.M. Langer & G.D. Hill. Cambridge University Press, 40 West 20th Street, New York, New York 10011. 1992. xiv. 387 pp. \$89.95 (hardcover); \$29.95 (paper). ISBN 0-521-40545-9 (hardcover); 0-521-40563-7 (paper).

This volume covers primary herbaceous species, and those most likely to be encountered in temperate and subtropical areas. The book has introductory chapters about human population growth and the importance of crop production, as well as information on plant morphology. These chapters are followed by summaries of the major plant families containing important agricultural crops. Within each family, discussions generally treat each genus separately. The final chapter is a discussion of physiological basis of crop yields.

Agricultural Policies in Developing Countries. Frank Ellis. Wye Studies in Agricultural and Rural Development. Cambridge University Press, 40 West 20th Street, New York, New York 10011. 1992. xvi. 357 pp. \$69.95 (hardcover); \$29.95 (paper). ISBN 0-521-40004-x (hardcover); 0-521-39584-4 (paper).

The entire first section of this book deals with analysis of agricultural policy. Factors such as markets, political system, social status, and concepts of policy analysis processes are treated there. Part two comprises descriptions of various policies affecting agriculture. Policies discussed include pricing, marketing, agricultural supports (input), credit, mechanization, land reform, research, and irrigation. The final section includes discussions of nonagricultural policies (such as women's issues and policies on food distribution) with significant impact on agricultural policy.

Anales del Instituto de Biología, Serie Botánica. Volume 62, Number 1. Fernando Chiang Cabrera (ed.). Universidad Nacional Autónoma de México, Coordinador de la Biblioteca del Instituto de Biología, Apartado Postal 70-233, 04510 México, D.F., México. 1991. 106 pp. Price unknown (paper). ISSN 0374-5511.

Five articles in this volume cover topics ranging from ethnobotanical studies, to phytochemical examinations, to physiological ecology, and classical taxonomy. Each article is accompanied by both a Spanish and English abstract. The articles themselves are in either Spanish or English.

Anales del Instituto de Biología, Serie Botánica. Volume 62, Number 2. Fernando Chiang Cabrera (ed.). Universidad Nacional Autónoma de México, Coordinador de la Biblioteca del Instituto de Biología, Apartado Postal 70-233, 04510 México, D.F., México. 1991. 180 pp. Price unknown (paper). ISSN 0374-5511.

Five articles in this volume cover topics ranging from sampling aflatoxins in peanut butter to floristic studies and descriptions of new species. Each article is accompanied by both a Spanish and English abstract. The articles themselves are in either Spanish or English.

Annual Review of Entomology, volume 37. Thomas E. Mittler, Frank J. Radovsky, & Vincent H. Resh. Annual Reviews Inc., 4139 El Camino Way, P.O. Box 10139, Palo Alto, California 94306. 1992. viii. 688 pp. \$44.00 [\$49.00 outside U.S.A.] (cloth). ISBN 0-8243-0137-4. ISSN 0066-4170.

Twentysix papers on topics ranging from labor allocations in insect societies, to mode of action of bacterial parasite toxins, to criminal forensic uses of entomology are found in this volume. Over fifty authors have contributed to these articles. Major themes (several papers each) are pest management and host/parasite relationships.

Biochemistry and Molecular Biology of Plant-Pathogen Interactions. C.J. Smith (ed.). Proceedings of the Phytochemical Society of Europe 32. Oxford Science Publications, Clarendon Press, Oxford University Press, Walton Street, Oxford OX2 6DP, Great Britain. Available in the United States from Oxford University Press, 200 Madison Avenue, New York, New York 10016. 1991. xvi. 291 pp. Price unknown (hardcover). ISBN 0-19-857734-6.

Seventeen papers produced by 59 authors comprise this volume. As might be expected from the series title, European researchers are well represented. Most of the papers deal with some aspect of molecular biology, with very few strictly biochemical or phytochemical in nature. Some of the articles treat specific host/pathogen problems, while others cover groups of pathogens and/or hosts.

The Biology of Vines. Francis E. Putz & Harold A. Mooney (eds.). Cambridge University Press, 40 West 20th Street, New York, New York 10011. 1992. xvi. 526 pp. \$120.00 (cloth). ISBN 0-521-39250-0.

Many of the eighteen papers (written by 25 authors) comprising this book were presented at a symposium on vines. Topics covered include anatomy and morphology, distribution and evolution, water transport, metabolism (primary and secondary), ecological attributes, reproduction, and economic importance.

Cuadernos del Instituto de Biología 13, Contribución a la Biología Mexicana por Helia Bravo Hollis: Una Guía Bibliográfica. Armando Butanda & Alfonso Delgado Salinas. Instituto de Biología, Universidad Nacional Autónoma de México, Apartado Postal 70-233, 04510, México, D.F., México. 1991. 44 pp. Price unknown (paper). ISBN 968-36-2110-4.

This volume includes a summary of the scientific activities (especially the botanical endeavors) of Helia Bravo Hollis. Doctor Hollis primarily studied cacti, but her interests also dealt with other plants, and botanical biographies, and protozoans. Her publications, new taxa, and taxa named in her honor are listed.

Endocytosis, Exocytosis and Vesicle Traffic in Plants. C.R. Hawes, J.O.D. Coleman, & D.E. Evans (eds.). Society for Experimental Biology, Seminar Series 45. Cambridge University Press, 40 West 20th Street, New York, New York 10011. 1992. xiv. 252 pp. \$85.00 (hardcover). ISBN 0-521-32844-6.

The volume presents twelve papers by a total of 32 authors. Papers treat ultrastructure, molecular characteristics, isolation procedures, and examination techniques of vesicles from various types of cells. The book concludes with a summary of possible future research in this field.

The Gene Civilization. François Gros. Translated from the French by Lee F. Scanlon. McGraw-Hill Horizons of Science Series, McGraw-Hill, Inc., 1221 Avenue of the Americas, New York, New York 10020. 1992. 136 pp. \$9.95 (paper). ISBN 0-07-024963-6.

This book as part of a series presenting science to the nonscientific public, deals with concepts of molecular biology, advances in biotechnology and the ethical issues that future advances in biotechnology and molecular biology may present.

Genetics and Conservation of Rare Plants. Donald A. Falk & Kent A. Holsinger (eds.). Oxford University Press, 200 Madison Avenue, New York, New York 10016. 1991. xviii. 283 pp. \$49.95 (cloth). ISBN 0-19-506429-1.

The core of this book consists of papers from a conference on rare plants held in March 1989. Additional papers were written specifically for the book and not included in the conference. A total of fourteen papers from 26 authors are included. Major themes of the book are population biology/genetics of rare plants, sampling of genetic diversity in rare plants, use of transported collections, and strategies for conserving genetic diversity.

Jeff Ball's 60-Minute Vegetable Garden, Just One Hour a Week for the Most Productive Vegetable Garden Possible. Jeff Ball. Collins Books, MacMillan Publishing Company, 866 Third Avenue, New York, New York 10022. 1992. xii. 228 pp. \$13.00 [\$16.95 in Canada] (paper). ISBN 0-02-030376-9.

A practical guide to home gardening, this book not only suggests activities for the gardener, but details how to accomplish the suggested activity. For example in the section on raised beds, the author not only discusses the pros and cons of raised beds, and suggests that they be used, but the book includes plans (with a shopping list, and information on tools needed, etc.) to construct frames for raised beds. Similar information appears for constructing trellises, compost bins, irrigation systems, and other items.

Life in the Universe. Jean Heidemann. Translated from the French by Isabel A. Leonard. McGraw-Hill Horizons of Science Series, McGraw-Hill, Inc., 1221 Avenue of the Americas, New York, New York 10020. 1992. 113 pp. \$9.95 (paper). ISBN 0-07-027887-3.

This book as part of a series presenting science to the nonscientific public, deals with cosmic evolution, processes leading to the existence of life, and prospects of extraterrestrial life (intelligent or otherwise).

Magic Gardens, A Modern Chronicle of Herbs and Savory Seeds. Rosetta E. Clarkson. Foreword by Sal Gilbertie. American Gardening Classics. Collier Books, MacMillan Publishing Company, 866 Third Avenue, New York, New York 10022. 1992. xxii. 369 pp. \$12.95 [\$16.95 in Canada] (paper). ISBN 0-02-030976-7.

Drawing from the publications of herbalists such as Gerard, Johnson, Matthioli, and Parkinson, this book contains information on cultivation and uses for many of the plants studied by these early authors. The book is organized around constructing gardens on a theme (pickling, scents, symbols, etc.).

The Manual of Cultivated Orchid Species, Third Edition. Helmut Bechtel, Phillip Cribb, & Edmund Launert. The MIT Press, 55 Hayward Street, Cambridge, Massachusetts 02142. 1992. 585 pp. \$85.00 (cloth). ISBN 0-262-02339-3.

The first section of this book contains general information about orchid taxonomy, ecology, and cultivation. Most of the text is devoted to alphabetically arranged descriptions of each genus and species of orchids known to be cultivated. Generic descriptions include morphological descriptions, distribution information, number of species, etymological information, taxonomy, type species listing, and cultivation information. Species descriptions contain morphological descriptions, distribution information, historical notes, and synonymy. The classification system of Dressler (*The Orchids, Natural History and Classification* [1981. Harvard University Press.] and modified in Telopea [2(4):413-424. 1983.]) is followed. The book contains over 140 pages of exquisite color plates.

New Flora of the British Isles. Clive Stace. Cambridge University Press, 40 West 20th Street, New York, New York 10011. 1992. xxx. 1226 pp. \$75.00 (flexible waterproof). ISBN 0-521-42793-2.

This book includes pteridophytes and gymnosperms as well as angiosperms. The text includes plants found on the outer islands (Outer Hebrides, Orkney, Shetlands, Scilly, and Channel islands) as well as the larger land masses of the British Isles. In addition to the native flora, Stace has included non-native species regularly encountered within the area (for instance, date palm is included since even though they are killed by the first frosts of the year, plants are regularly encountered as seedlings in garbage piles). This approach has added to the length of the book, but significantly improved its usefulness for those not familiar with the flora and which species might be native as opposed to non-native. The book has a cover designed for field use, but its thickness makes it rather bulky for a pocket.

The New Perennials Preferred. Helen Van Pelt Wilson. Foreword by Elvin McDonald. American Gardening Classics. Collier Books, MacMillan Publishing Company, 866 Third Avenue, New York, New York 10022. 1992. xx. 315 pp. \$15.95 [\$19.95 in Canada] (paper). ISBN 0-02-082661-3.

The first section of the book is devoted to general information on gardening, and cultivating perennials in particular. Most of the text comprises a seasonally arranged discussion of cultivated (primarily herbaceous) perennials. The seasonal discussion includes a substantial treatment of plants attractive during the winter. Lists of plants suitable for specific purposes (fragrance production, shade tolerance, etc.) are also included.

Nitrogen Metabolism of Plants. Proceedings of the Phytochemical Society of Europe, 33. K. Mengel & D.J. Pilbeam (eds.). Oxford Science Publications, Oxford University Press, 200 Madison Avenue, New York, New York 10016. 1992. xii. 289 pp. \$89.00 (hardcover). ISBN 0-19-857752-4.

Fortynine researchers from Canada and the United States, as well as Europe, have contributed sixteen papers to this volume. Most of the papers deal with nitrogen fixation or transport, with other papers on metabolism of nitrogen, and nitrogen based plant toxins.

Our Changing Climate. Robert Kandel. Translated from the French by Nicholas Hartmann. McGraw-Hill Horizons of Science Series, McGraw-Hill, Inc., 1221 Avenue of the Americas, New York, New York 10020. 1992. 126 pp. \$9.95 (paper). ISBN 0-07-033710-1.

This book as part of a series presenting science to the nonscientific public, deals with general concepts of climate, models for study of climate, prospects of the greenhouse effect and ozone depletion, and political influences on climate research.

Physiology of Trees. A.S. Raghavendra (ed.). John Wiley & Sons, Inc., 1 Wiley Drive, Somerset, New Jersey 08875. 1991. xii. 509 pp. \$110.00 (cloth). ISBN 0-471-50110-7.

Thirtythree authors contributed twenty papers to form this book. Topics include photosynthesis and respiration, photosynthate distribution, nitrogen fixation, water relations, seedling growth, wood formation, periodicity, flowering, growth forms, abscission, responses to stress, tissues culture, use of growth regulators, physiological modeling, and wound responses.

Practical Taxonomic Computing. Richard J. Pankhurst. Cambridge University Press, 40 West 20th Street, New York, New York 10011. 1992. xii. 202 pp. \$49.95 (hardcover). ISBN 0-521-41760-0.

After some brief introductory material, this book gets to the meat of the issues of taxonomic computing with discussions of databases, principles of classification, conventional identification procedures, identification procedures facilitated by computer, and sample applications of computer identification. The final chapter includes a discussion of expert systems technology and its potential impact on taxonomic computing. An appendix contains interesting mathematical derivations of formulae for estimating usefulness of taxonomic characters.

Sweet Potato An Untapped Food Resource. Jennifer A. Woolfe. Published in collaboration with the International Potato Center, Lima Perú, by the Cambridge University Press, 40 West 20th Street, New York, New York 10011. 1992. xvi. 643 pp. \$130.00 (hardcover). ISBN 0-521-40295-6.

This comprehensive summary on use of sweet potato for human and livestock consumption includes chapters on history of cultivation, chemical constituents, nutritive value, toxicity, storage, processing, cooking, livestock utilization, and human consumption methods and histories. An appendix contains recipes for sweet potato dishes from around the world.

Vegetation of New Zealand. Peter Wardle. Cambridge University Press, 40 West 20th Street, New York, New York 10011. 1991. xx. 672 pp. Price unknown (cloth). ISBN 0-521-25873-1.

A comprehensive treatment of the vegetation of this island nation. The book includes sections on the biotic and abiotic environment, origins and history of the flora, growth forms in various habitats, reproductive processes, and a discussion of a vegetation classification system. The classification system is essentially hierarchical, with major vegetation types (forest, grassland, etc.) broken into species or region specific subunits. Separate discussions of some of the outlying islands are included.

Yardening, The Nongardener's Guide to Creating a Beautiful Landscape. Jeff & Liz Ball. MacMillan Publishing Company, 866 Third Avenue, New York, New York 10022. 1992. xiv. 268 pp. \$24.95 [\$32.50 in Canada] (cloth). ISBN 0-02-506431-2.

This book, intended for persons with an interest in taking care of their garden (yarden), but without extensive experience or time, deals with topics ranging from taking care of the grass to growing flowers and vegetables. The information is presented in a readily accessible fashion with references for further details.

INDEX TO AUTHORS, VOLUME 73

- Adler, L. 155
Benítez de Rojas, C. 270
Brown, G.K. 57
Carlquist, S. 98
Clements, M.A. 117
Dudley, T.R. 169
Fernández N., R. 435
Grayum, M.H. 30
Hernández S., L. 429
Hoch, P.C. 456, 460
Holmes, W.C. 159
Jones, D.L. 3, 117
Jones, S.D. 274, 381
Keil, D.J. 57, 259, 312
Lott, E.J. 277
Lowrie, A. 98
MacRoberts, B.R. 49, 439
MacRoberts, M.H. 49, 439
Martínez, M. 270
McLaughlin, S.P. 353
McNeal, D.W. 307
Nesom, G.L. 61, 119, 137, 140, 186,
203, 247, 264, 267, 318, 321,
326, 330, 338, 384, 389, 416,
425, 429, 463
Ochoa, C.M. 166, 180, 183, 378
Panero, J. 143
Raven, P.H. 456
Rayner, T.G.J. 77, 445
Reveal, J.L. 345
Robinson, H. 65, 149
Santamour, Jr., F.S. 169
Semple, J.C. 449
Skinner, S. 155
Turner, B.L. 1, 7, 14, 16, 18, 24, 27,
40, 59, 143, 242, 251, 255,
261, 281, 302, 304, 348, 350
Waksman, N. 435
Wiehler, H. 220
Windler, D. 155
Wipff, J.K. 274, 381

INDEX TO TAXA, VOLUME 73

- Abies* 44, 120, 138
Abildgaardia 383
Abutilon 371
 californicum 371
 parvulum 371
 sonorae 371
Acacia 333, 360, 363, 370
 angustissima 370
 greggii 360, 363, 370
Acalypha 370
 neomexicana 370
 ostryaefolia 370
Acanthaceae 140, 142, 366
Acer 51, 120
 rubrum 51
Aceraceae 51
Achimenes 220
 admirabilis 220
 erecta 220
Acourtia 367
 nana 367
 thurberi 367
 wrightii 367
Adiantaceae 366
Agalinus 52
 obtusifolia 52
Agastache 358
 wrightii 358
Agavaceae 354, 360, 374
Agave 141, 333, 359, 360, 365, 374
 palmeri 360, 374
 parviflora 359, 365, 374
 schottii 374
 var. *schottii* 374
Ageratina 304-306
 Ageratina (cont.)
 chimalapana 304-306
 cruzii 304, 305
 etlensis 304, 305
 kochiana 304, 305, 306
 kochii 306
 ligustrina 305
 mairetiana 306
 subg. *Neogreenella* 304
 seleri 306
Agropyron 363, 375
 trachycaulum 363, 375
Aizoaceae 366
Alepidocline 246, 258
Aletris 51
 aurea 51
Allionia 371
 incarnata 371
Allium 307-310, 362, 374
 bolanderi 307-309
 var. *bolanderi* 308, 309
 var. *mirabile* 307-309
 dichlamydeum 310
 macropetalum 362, 374
 mirabile 307
 peninsulare 307, 310
 var. *franciscanum* 307, 310
 var. *peninsulare* 310
Allophyllum 372
 gilioides 372
Alnus 404
Alopecurus 359, 375
 carolinianus 359, 375
Aloysia 358, 359, 374
 gratissima 358, 374

- Aloysia* (cont.)
 wrightii 359, 374
Altamirania 145
 pachyphylla 145
Amaryllidaceae 51
Amaranthaceae 366
Amaranthus 366
 albus 366
 palmeri 366
 torreyi 366
Ambrosia 363, 367
 aptera 367
 confertiflora 363, 367
 psilostachya 363, 367
 trifida 363, 367
Amelanchier 124, 129, 134
 alnifolia 124, 129
 canadensis 129
 var. *pumila* 129
 pumila 124, 129
Amellus 455
 villosus 455
Amoreuzia 369
 palmatifida 369
Amorphae 428
Ampelopsis 132
 quinquefolia 132
 var. *heptaphylla* 132
 var. *pubescens* 132
Amphicarpaea 285, 291, 301
 angustifolia 285
 ovalifolium 291
 pulchella 291
Amsinckia 368
 intermedia 368
Amsonia 365
 grandiflora 365
Anacardiaceae 51
Andropogon 51, 55
 gerardii 55
 ternarius 51
Anemone 373
 tuberosa 373
 Anemopsis 373
 californica 373
Anisacanthus 366
 thurberi 366
Androsace 372
 occidentalis 372
Anoda 371
 abutiloides 371
 cristata 371
Anthemideae 422
Anthericum 359, 374
 torreyi 359, 374
Antirrhinum 373
 nuttallianum 373
Aphanostephus 13
Apiaceae 51, 366
Apocynaceae 179, 367
Apodanthera 369
 undulata 369
Appendicula 117, 118
 australiensis 117, 118
Aquifoliaceae 51
Arabis 368
 perennans 368
Araceae 30
Arbutus 268, 404
Arctostaphylos 267
Argemone 372
 pleiacantha 372
 ssp. *pleiacantha* 372
Argythamnia 370
 neomezicana 370
Aristida 51, 275, 356, 360, 362, 375
 adscensionis 375
 californica 362
 hamulosa 362, 375
 orcuttiana 375
 purpurea 360, 362, 375
 var. *longiseta* 360, 362, 375
 var. *purpurea* 375
 ternipes 360, 362, 375
 virgata 51
 wrightii 362

- Aristolochiaceae 367
Aristolochia 367
 watsoni 367
Aronia 124, 129, 130
 arbutifolia 124, 129, 130
 alnifolia 129
Artemisia 367
 ludoviciana 367
 Asclepiadaceae 277, 280, 367
Asclepias 356, 367
 asperula 367
 ssp. *capricornu* 367
 brachystephana 367
 involucrata 367
 nyctaginifolia 367
 subverticillata 367
Aspilia 145
 pachyphylla 145
Aspiliopsis 145
 pachyphylla 145
Aster 51, 55, 187, 210, 338, 339,
 359, 367, 421, 422
 dumosus 51, 55
 ericoides 55
 falcatus 359, 367
 ssp. *commutatus* 359, 367
 sect. *Oritrophium* 338
 spinosus 187
 subulatus 367
 var. *ligulatus* 367
 Asteraceae 1, 2, 7, 14-16, 18, 24, 27,
 29, 40, 48, 51, 57-59, 65, 75,
 76, 119, 123, 143, 148, 149,
 154, 186, 200-203, 219, 242,
 246, 255, 258-261, 264, 266,
 267, 269, 302, 304, 306, 312,
 317, 318, 320, 326, 329, 338,
 344, 348, 350, 356, 357, 367,
 416, 420-424, 463
 Astereae 57, 58, 119, 122, 123, 186,
 200-203, 219, 264, 266, 267,
 269, 318, 320, 326, 329, 338-
 340, 343, 344, 416, 418, 420,
 Astereae (cont.) 422, 423, 449, 455,
 463
Astragalus 357, 359, 370
 allochrous 370
 arizonicus 370
 nothozys 370
 nuttallianus 359, 370
 var. *cedroensis* 359, 370
 var. *nuttallianus* 370
 wootoni 370
Athenaea 272
 glabra 272
 sylvarum 272
Atriplex 218, 369
 canescens 369
 elegans 369
 ssp. *elegans* 369
Avena 375
 fatua 375
Ayenia 373
 compacta 373
Aynia 75
Azolla 364, 366
 mexicana 364, 366
 Azollaceae 366
 Baccharidinae 339
Baccharis 65, 66, 69, 72, 357, 367
 brachyphylla 367
 brasiliana 65, 66, 69
 glutinosa 367
 montevidensis 72
 pteronioides 367
 sarothroides 367
 thesioides 367
Bahia 367
 absinthifolia 367
 var. *dealbata* 367
Baileya 367
 multiradiata 367
Bambusa 436
Benthamidia 178
 florida 178

Benthamia florida (cont.)var. *urbiniana* 178*Berberidaceae* 368*Berberis* 368 *haematocarpa* 368*Berlandiera* 360, 367 *lyrata* 360, 367 var. *macrophylla* 367*Berula* 364, 366 *erecta* 364, 366*Bidens* 77-88, 90, 94-97, 363, 364,

367, 445, 447, 448, 509

acuticaulis 82 *abyssinica* 80 var. *glabrata* 80 *aurea* 363, 367 *bequaertii* 81 *bigelovii* 367 *camporum* 88 *crocea* 81, 82 *diversa* 77, 82 *ferulaefolia* 367 *flabellata* 82 *grantii* 88 *holstii* 77, 90 *kefensis* 445, 447 *kivuensis* 83, 86 *kotschyi* 83, 84, 88 *laevis* 364, 367 *leptocephala* 367 *leptolepis* 84 *negriana* 95 *palustris* 84 *paupercula* 85, 87, 509 var. *filirostris* 509 *praecox* 85 *rubicundula* 85 *rupestris* 85 *schimperi* 78 *schlechteri* 86 *setigera* 86, 88 ssp. *bipinnato-partita* 88 var. *lobata* 86*Bidens setigera* (cont.) ssp. *setigera* 88 *somaliensis* 86, 87 *steppia* 77, 87, 94 var. *ambacensis* 87 var. *elskensis* 87 var. *leptocarpa* 87 var. *steppia* 87 *straminoides* 87 *stuhlmannii* 79 *taylorii* 86 *ugandensis* 445, 448 *urceolata* 84, 85*Bignoniaceae* 260, 368*Bocconia* 20*Boerhaavia* 356, 362, 371 *coccinea* 371 *coulteri* 371 *erecta* 371 *gracillima* 371 *intermedia* 371 *spicata* 371*Boisduvalia* 456-459 *cleistogama* 458 *concinna* 457 sect. *Currantia* 458 *densiflora* 457 *douglasii* 457 *glabella* 459 *pallida* 458 *macrantha* 458 *pygmaea* 458 *stricta* 458 *subulata* 457 *torreyi* 458*Boraginaceae* 357, 368*Bothriochloa* 360, 363, 375 *barbinodis* 360, 375 *saccharoides* 363, 375*Bouteloua* 356, 359, 360, 362, 375 *aristidoides* 362, 375 *barbata* 375 *chondrosioides* 360, 375

Bouteloua (cont.)*curtipendula* 360, 362, 375*eludens* 359, 375*eriopoda* 360, 362, 375*gracilis* 375*hirsuta* 360, 375*parryi* 362*radicosa* 362*repens* 360, 362, 375*rothrockii* 362, 375*Bowlesia* 366*incana* 366*Brachiaria* 375*arizonica* 375*Brahea* 438*dulcis* 438*nitida* 438

Brassicaceae 357, 368

Brickellia 348, 349, 356, 358, 367*baccharidea* 367*californica* 358, 367*chlorolepis* 367*coulteri* 367*floribunda* 358, 367*gentryi* 348*lanata* 348, 349*venosa* 367*Bromus* 375*arizonicus* 375*catharticus* 375*marginatus* 375*rubens* 375*Bulbostylis* 374, 381-383*barbata* 381*capillaris* 374, 381, 382*ciliatifolia* 381, 382var. *coarctata* 381, 382*juncoides* 381, 382*Bumelia* 373*lanuginosa* 373*Burmannia* 51*capitata* 51

Burmanniaceae 51

Cacalia 44, 68, 73, 74*pachyphylla* 44*praecox* 74*rigiophylla* 73

Cactaceae 354, 357, 368

Calandrinia 372*ciliata* 372*Calea* 143, 145, 148*pachyphylla* 145*skutchii* 143, 148*standleyi* 143, 145*Calliandra* 360, 370*eriophylla* 360, 370*humilis* 370var. *reticulata* 370*Calochortus* 374*ambiguus* 374*kennedyi* 374var. *munzii* 374*Calopogon* 51*tuberosus* 51*Calycoseris* 367*wrightii* 367*Calyptridium* 372*monandrum* 372

Campanulaceae 52, 369

Camissonia 372*californica* 372*chamaenerioides* 372

Capparidaceae 369

Caprifoliaceae 52, 369

Capsella 368*bursa-pastoris* 368

Carduales 75

Cardueae 312, 317

Carduus 313*breweri* 313*Carex* 363, 374, 382*chihuahuensis* 363, 374*longii* 382*praegracilis* 374*Carlwrightia* 366*arizonica* 366

- Carminatia* 367
 tenuiflora 367
Carnegiea 368
 gigantea 368
Caryophyllaceae 369
Carya 120
Castilleja 137-139, 247-250, 384-415
 albobarbata 414
 altorum 409, 411
 arvensis 385
 aspera 138
 aurea 410
 auriculata 389-391, 393, 397-401,
 404, 413, 414
 var. *auriculata* 389, 397-400,
 404, 413
 var. *verecunda* 389, 398, **399**,
 400, 414
 bella 249
 canescens 391
 sect. *Castilleja* 390, 400, 404, 407,
 409, 410
 subg. *Castilleja* 390, 414
 chiapensis 395-397, 412
 chlorosceptron 137, 138, 385,
 387
 conzattii 248, 249
 cryptandra 384, 386, 387
 ctenodonta 409, 411
 dendridion 248
 durangensis 137, 138
 sect. *Epichroma* 390, 409, 410
 sect. *Euchroma* 138, 247, 248,
 384, 387, 388, 409
 falcata 248, 249, 386
 filiflora 389, 406, 407, 408, 412
 fissifolia 389, 390
 franciscana 390, 409
 gracilis 410
 sect. *Hemichroma* 389
 hirsuta 248, 386
 integra 386
 integrifolia 385, 389, 390,
 Castilleja integrifolia (cont.) 394-
 397, 404, 412
 var. *alpigena* 395
 jepsonii 390
 jilquilpana 250
 laza 391, 393
 linariifolia 390, 410
 sect. *Linariifolia* 389
 linariifolia 410
 linifolia 407, 411
 longibracteata 395
 longiflora 391, 397, 399
 macrostigma 249
 macvaughii 411
 moranensis 249
 nervata 386, 387
 nitricola 385
 orizabae 409
 ortegae 410
 papilionacea 384, **385**, 386
 patriotica 390, 411
 pectinata 395, 409, 411
 perelegans 389, 396, 408, 412
 pterocaulon 410
 purpusii 409, 411
 porphyrosceptron **384**
 retrorsa 393
 rhizomata 400, 401, 407, 413
 roei 410
 saltensis 250
 scabridula 391, 393
 scorzoneraefolia 138, 248, 385-
 387, 409
 setosa 393
 sphaerostigma 249, 250
 stipifolia 389, 404, 405-407, 412
 subalpina 409
 subinclusa 389, 390, 409, 411
 subsp. *franciscana* 409
 var. *franciscana* 389, **409**, 411
 tancitaroana 389, 393, 397, 401-
 403, 413
 tapeinoclada 407, 408, 412

Castilleja (cont.)*tayloriorum* 390*tenuiflora* 385, 386, 389-395, 397,
399, 401, 404, 405, 407, 408,
413var. *tenuiflora* 391-394, 401,
413var. *verecunda* 401var. *xylorrhiza* 389, 392, **393**,
394, 413*tenuifolia* 410*tokucensis* 247-249*venusta* 410*wootonii* 390*xylorrhiza* 393*zempoaltepetlensis* 247-249,
387*Castillejinae* 414*Caulanthus* 368*lasiophyllus* 368*Ceanothus* 431*Celmisia* 338-340, 343, 344*Celtis* 358, 373*pallida* 373*reticulata* 358, 373*Cenchrus* 375*insertus* 375*Ceratophyllaceae* 369*Ceratophyllum* 364, 369*demersum* 364, 369*Cercidium* 370*floridum* 370*Cercocarpus* 213*Chaenactis* 367*stevioides* 367*Chaetopappa* 416-420, 422, 423*asteroides* 418, 419*bellidiflora* 419*bellidifolia* 419*bellioides* 419, 420*effusa* 418-420*elegans* 416, 417, 420*ericoides* 417, 419, 420*Chaetopappa* (cont.)*hersheyi* 417-419*imberbis* 419*keerlioides* 419*parryi* 419*plomoensis* 419*pulchella* 419, 420*Chamaecrista* 370*nictitans* 370*Chamaesyce* 356, 362, 363, 370*albo-marginata* 370*arizonica* 370*capitellata* 370*florida* 370*hirta* 370*hyssopifolia* 370*melanadenia* 370*pediculifera* 370*serpyllifolia* 370*setiloba* 370*vermiculata* 363, 370*Chaptalia* 441*Cheilanthes* 359, 366*lindheimeri* 366*wootoni* 366*wrightii* 366*Chelidoniaeae* 330*Chelidonioideae* 331*Chenopodiaceae* 369*Chenopodium* 369*berlandieri* 369var. *sinuata* 369*desiccatum* 369var. *leptophylloides* 369*incanum* 369*Chiliotrichum* 339*Chilopsis* 358, 368*linearis* 358, 368*Chloracantha* 187, 201, 202*spinosa* 202*Chlorideae* 274*Chloris* 362, 375*crinita* 375

Chloris (cont.)*virgata* 375*Choenomeles* 124, 130*japonica* 124, 130*Chorizanthe* 345, 346*virgata* 345, 346*Chrysocoma* 70, 73*arborea* 73*crassa* 70*cymosa* 70*phosphorica* 73*Chrysolaena* 75*Chrysopsis* 187, 449-451, 453*bolanderi* 449*camphorata* 451*echioides* 450*fastigiata* 451*hirtella* 197*villosa* 449-451, 453var. *bolanderi* 449var. *camphorata* 451var. *echioides* 450var. *fastigiata* 451var. *scabra* 453*viscida* 453subsp. *cinerascens* 453*Cirsium* 312-316, 360, 367*breweri* 312, 313var. *breweri* 313var. *canescens* 312, 313*californicum* 314, 315, 317*campylon* 313*coulteri* 314sect. *Dermatolepis* 313, 316*douglasii* 312, 313var. *breweri* 312, 313var. *canescens* 312, 313var. *douglasii* 312, 313*fontinale* 312, 313var. *campylon* 312, 313var. *fontinale* 313var. *obispoense* 313*neomexicanum* 367*Cirsium* (cont.)*occidentale* 312-315, 317var. *californicum* 312, 315var. *candidissimum* 314, 315var. *compactum* 315var. *coulteri* 313, 314var. *occidentale* 313-315var. *venustum* 313-315*ochrocentrum* 360, 367*pastoris* 314*proteanum* 314*Clematis* 373*drummondii* 373*Clibadium* 149-154*alatum* 149, 151*divaricatum* 149*funkiae* 149*glabrescens* 153*laxum* 151*napoense* 149, 152*pentaneuron* 151*rimachii* 149*surinamense* 149, 150, 154*vargasianum* 149*zakii* 149, 153*Clitoria* 291*broussonettii* 291

Clusiaceae 52

Cnicus 313*breweri* 313*Cocculus* 358, 371*diversifolius* 358, 371

Cochlospermaceae 369

Codonanthe 220, 221*crassifolia* 221*erubescens* 221*Cologania* 281-301*affinis* 291*angustifolia* 282-285, 287-289, 294var. *stricta* 288*australis* 293*biloba* 282, 284, 289, 290, 291,

295, 296, 300

Cologania (cont.)

- broussonettii* 281-283, 285, 288,
291-294, 298, 300
 - capitata* 282, 284, 294, 297
 - confusa* 288
 - congesta* 293
 - cordata* 282, 284, 294, 297, 300
 - deamii* 298
 - erecta* 298
 - glabrior* 293
 - grandiflora* 289, 291, 293
 - hintoniorum* 281, 284, 286, 290,
291, **295**, 296
 - hirta* 282, 284, 290, 295, 296
 - houghii* 298
 - humifusa* 296
 - humilis* 298
 - intermedia* 285, 288
 - jaliscana* 282, 288, 293
 - lemmonii* 296
 - longifolia* 285, 288
 - var. stricta* 288
 - lozanii* 293
 - martia* 285
 - mezicana* 285
 - nelsonii* 289
 - obovata* 282-284, 289, 296, 297
 - ovalifolia* 281, 282, 291, 293, 294
 - pallida* 281, 282, 284, 288, 292,
298
 - pringlei* 288, 293, 296
 - procumbens* 281, 282, 284, 294,
298-300
 - pulchella* 291, 300
 - var. racemosa* 300
 - purpurea* 289
 - racemosa* 284, 290, 291, 296, 300
 - rufescens* 293
 - tenuis* 293
- Columnnea* 220, 222, 223
- flipendula* 222**
 - glicensteinii* 223**
 - nicaraguensis* 223

Commelina 374*erecta* 374

Commelinaceae 374

Commicarpus 371*scandens* 371

Compositae 13, 17, 23, 48, 75-77,
96, 97, 122, 148, 159, 200-
202, 219, 257, 258, 260, 343,
344, 421-423, 445, 448, 449,
455

Condalia 373*mezicana* 373*spathulata* 373

Convolvulaceae 357, 369

Convolvulus 369*arvensis* 369*equitans* 369*Conyza* 74, 187, 201, 367*canadensis* 367*coulteri* 367*stellata* 74*Coreocarpus* 359, 367*arizonicus* 359, 367

Coreopsis 7-13, 27-29, 51, 77, 79,
80, 88-96, 509

abyssinica 88*var. bipinnato-partita* 88*badia* 89*baumii* 89*crataegifolia* 89sect. *Electra* 13*ezaristata* 90*galeottii* 11***guanajuatensis* 27-29***holstii* 77, 90*kilimandscharica* 90, 91*leptoglossa* 91*lineata* 91*linifolia* 51*lupulina* 92*macvaughii* 27, 29*mezicana* 9-11*forma hyperdasya* 11

Coreopsis mexicana (cont.)

- var. *hyperdasya* 10, 11
- var. *mexicana* 9
- morotonensis* 509
- mutica* 7-13
 - var. *carnosifolia* 8, 9, 11-13
 - var. *guerreroana* 7, 8, 11, 12
 - var. *holotricha* 8, 9, 11, 12
 - var. *leptomera* 8-10, 11
 - var. *mexicana* 10, 11
 - var. *microcephala* 8, 9, 11-13
 - var. *multiligulata* 8, 11-13
 - var. *mutica* 7-12
 - var. *simplicifolia* 7, 9-11
 - var. *subvillosa* 7-9, 11, 12

ochracea 92*odora* 509*pinnatipartita* 93sect. *Pseudoagarista* 13, 27, 29*queretarensis* 27, 29*rudis* 29*scabrifolia* 93*schimperii* 95*scopulorum* 509*steppia* 77, 94

Cornaceae 169, 179

Cornus 120, 169-179*disciflora* 177var. *floccosa* 177*florida* 169-179

cv. 'Barton' 175

cv. 'Cloud 9,' 175

subsp. *florida* 175, 176var. *florida* 174var. *pringlei* 169, 171, 174, 178,
179subsp. *urbiniana* 169, 170, 172-
178var. *urbiniana* 173, 174, 178*pringlei* 171-174, 178*urbiniana* 173, 178*Cortaderia* 378*Corydalis* 372*Corydalis* (cont.)*aurea* 372*Coryphantha* 365, 368*recurvata* 365*scheeri* 365, 368var. *robustispina* 365, 368*vivipara* 368var. *bisbeeana* 368*Cottea* 375*pappophoroides* 375*Crassula* 369*erecta* 369

Crassulaceae 369

Crotalaria 155-158, 370*incana* 155-158var. *australis* 155var. *grandiflora* 155, 157var. *incana* 155, 157forma *microphylla* 156var. *nicaraguensis* 155*pumila* 370*Croton* 370*pottsii* 370*Cryptantha* 357, 368*angustifolia* 368*barbigera* 368*micrantha* 368*nevadensis* 368*pterocarya* 368*Otenium* 439-442*aromaticum* 439, 442*Cucurbita* 369*digitata* 369*foetidissima* 369

Cucurbitaceae 369

Cuscuta 369*erosa* 369*Cymophora* 242-246*hintonii* 243, 245*luckowana* 242-245*pringlei* 242, 243, 245*venezuelensis* 243*Cynanchum* 359, 366, 367

Cynanchum (cont.)*arizonicum* 359, 367*wigginsii* 366*Cynoctonum* 52*sessilifolium* 52*Cynodon* 275, 375*dactylon* 275, 375*Cyperaceae* 51, 124, 125, 135, 357,

374, 381, 383

Cyperus 356, 364, 374, 382*albomarginatus* 374*aristatus* 374*compressus* 382*dipsaceus* 374*esculentus* 374*flavicomus* 374*flavus* 374*haspan* 382*mutisii* 374*odoratus* 364, 374*pallidicolor* 374*polystachyos* 382*rusbyi* 374*sphaerolepis* 374*squarrosus* 374*wrightii* 374*Cytocymura* 75*Dactyloctenium* 382*aegyptium* 382*Dalbergaria* 220, 223-228*asteroloma* 225*crassa* 225*evolvens* 223*filifera* 224*lanata* 225*madisonii* 224*mastersonii* 225*medicinalis* 226*ornata* 227*picta* 226*polyantha* 227*schimpffii* 228*Dalbergaria* (cont.)*tutunendana* 228*Dalea* 370, 425-428*eriophylla* 425-428var. *eriophylla* 425, 427var. *frankeniioides* 425, 427var. *uniflora* 425, 427series *Eriophyllae* 425*nana* 370var. *carnescens* 370sect. *Parosela* 425*pogonathera* 370*pulchra* 370*uniflora* 425-427*Damnomenia* 339, 340, 344*Dasyliirion* 120, 267, 360, 374, 438*wheeleri* 360, 374*Datura* 373*meteloides* 373*Daucus* 366*pusillus* 366*Deiregyne* 128*confusa* 128*durangensis* 128*Delphinium* 373*scaposum* 373*Dendrobium* 3-6*finniganense* 3-6*adae* 5, 6*fleckeri* 3, 5, 6*Dendromecon* 330, 331, 333, 336*Dennstaedtiaceae* 51*Deprea* 270-273*cardenasiana* 273*glabra* 272*granulosa* 273*orinocensis* 272*paneroi* 270, 271, 273*sylvarum* 272*Descurainia* 368*pinnata* 368ssp. *halictorum* 368*sophia* 368

- Desmanthus* 370
 cooleyi 370
Desmodium 358, 370
 batocaulon 358, 370
Dicanthelium 51
 acuminatum 51
Dichelostemma 374
 pulchellum 374
Dicotyledonae 366
Digitaria 359, 360, 362, 375
 californica 360, 362, 375
 cognata 375
 insularis 359, 375
 sanguinalis 375
Diodia 55, 373
 teres 373
 virginiana 55
Diplostephium 339
Distephanus 76
Dodonaea 373
 viscosa 373
 var. *angustifolia* 373
Draba 368
 cuneifolia 368
 var. *integrifolia* 368
Drosera 52, 98-116, 441
 barbigera 99, 107
 bicolor 98, 109, 111, 112
 brevifolia 52
 capillaris 52
 citrina 98, 99-101, 105
 dichrosepala 102
 enodes 109
 sect. *Ergalium* 111
 subg. *Ergalium* 111
 sect. *Lamprolepis* 100, 102, 105,
 107, 109
 lasiantha 98, 100, 102-104
 macrophylla 98, 111, 113, 114
 subsp. *macrophylla* 111
 subsp. *monantha* 98, 111, 113,
 114
 nivea 98, 100, 104-106
Drosera (cont.)
 parvula 98
 peltata 111
 pycnoblata 100
 rechingeri 100
 subg. *Rorella* 100, 102, 105, 107,
 109
 salina 111
 scorpioides 102
 silvicola 98, 105, 107, 108
 stelliflora 98, 107, 109, 110
 whittakeri 98, 113, 115
 subsp. *aberrans* 98, 113, 115
 subsp. *whittakeri* 113
Droseraceae 52, 98
Drymonia 220, 229-231
 chiribogana 229, 230
 killipii 229
 punctulata 230
 stenophylla 231
 uninerva 231
Dryopetalon 359, 368
 runcinatum 359, 368
Duranta 378, 380
 rupestris 378, 380
Dyssodia 367
 pentachaeta 367
Echinocereus 368
 fasciculatus 368
 var. *fasciculatus* 368
 fendleri 368
 pectinatus 368
 var. *rigidissimus* 368
Echinochloa 375
 crusgalli 375
Echinopepon 369
 wrightii 369
Echinocoryne 75
Eirmocephala 75
Electra 9
 mexicana 9
Eleocharis 51, 125, 364, 374

Eleocharis (cont.)

- bella* 364, 374
- confervoides* 125
- montevidensis* 374
- palustris* 374
- tortilis* 51

Eleusine 382

- indica* 382

Elymus 375

- elymoides* 375
- triticoideus* 375

Elyonurus 375

- barbiculmis* 375

Encelia 367

- farinosa* 367

Enneapogon 375

- desvauzii* 375

Ephedra 366

- trifurca* 366

Ephedraceae 366*Epilobieae* 456, 459*Epilobium* 124, 127, 128, 134, 135, 372, 456-462

- Alpinae group 461

- sect. *Boisduvalia* 457

- brachycarpum* 124, 127, 128

- canum* 372

- sect. *Chamaenerion* 456

- ciliatum* 127

- cleistogamum* 458

- concinnum* 457

- sect. *Currantia* 458

- densiflorum* 457

- sect. *Epilobium* 460

- glabellum* 459

- howellii* 460, 461

- macranthum* 458

- oregonense* 461

- pallidum* 458

- paniculatum* 124, 127, 128

- ssp. *latifolium* 372

- pygmaeum* 458

- strictum* 458

Epilobium (cont.)

- subulatum* 457

- torreyi* 458

- sect. *Zauschneria* 456

Eragrostis 51, 275, 356, 360, 375, 377

- curvula* 375

- var. *conferta* 375

- chloromelas* 375

- cilianensis* 375

- echinocloidea* 375

- intermedia* 360, 375

- lehmanniana* 360, 375, 377

- pectinacea* 375

- var. *miserrima* 375

- var. *pectinacea* 375

- spectabilis* 51

Eremosis 68*Eria* 118

- australiensis* 118

Eriastrum 372

- diffusum* 372

Ericaceae 52*Ericameria* 359, 367

- laricifolia* 359, 367

Erigeron 119-123, 186-219, 262, 338, 342, 344, 359, 367, 416, 420, 422, 463

- accedens* 188, 189

- algidus* 187, 206, 209

- aphanactis* 186, 193, 194, 199, 205, 209

- var. *aphanactis* 193, 209

- var. *congestus* 193, 209

- argentatus* 208, 209

- asperugineus* 207, 210

- sect. *Asteroidea* 199

- aureus* 197

- austinae* 195, 198, 205, 211, 463

- barbellulatus* 199

- basaseachensis* 262

- bloomeri* 199, 205, 210

- var. *bloomeri* 210

Erigeron bloomeri (cont.)

- var. *nudatus* 210
- var. *pubens* 210
- breweri* 206, 210, 215
 - var. *breweri* 210
 - var. *porphyreticus* 210, 211
- californicus* 188, 189
- cavernensis* 218
- chiangii* 119-122
 - var. *chiangii* 120-122
 - var. *lamprocaulis* 119-122
- chrysopsidis* 186, 195-199, 211, 463
 - subsp. *austinae* 195, 198, 211, 463
 - var. *austinae* 186, 195, 196, 198, 211, 463
 - var. *brevifolius* 195-199, 463
 - subsp. *chrysopsidis* 195, 198, 211
 - var. *chrysopsidis* 195, 196, 198, 463
- clokeyi* 207, 211
- compactus* 199, 207, 211
- compositus* 187, 201, 205, 212
 - var. *compositus* 187
 - var. *discoideus* 187, 212
 - var. *glabratus* 187, 212
- concinus* 192, 193, 200, 206, 209, 212
 - var. *aphanactis* 193, 209
 - var. *concinus* 212
 - var. *condensatus* 212
- congestus* 193
- consimilis* 199, 211
- coronarius* 201
- corymbosus* 199
- curvifolius* 197
- coulteri* 204, 213
- divergens* 186, 188, 189, 206, 213, 214, 367
- eatonii* 186, 190-192, 202, 208, 213, 214

Erigeron eatonii (cont.)

- var. *eatonii* 213
- var. *nevadincola* 186, 191, 192, 213, 214
- var. *plantagineus* 191, 192
- var. *sonnei* 186, 190, 191, 213, 214
- elegantulus* 199
- filifolius* 207, 214
 - var. *filifolius* 214
 - var. *robustior* 214
- flagellaris* 207, 214
- flezuosus* 192
- foliosus* 122, 187
- sect. *Geniculactis* 199, 201
- hirtellus* 197
- inornatus* 205, 215
 - var. *inornatus* 215
- jonesii* 207, 215
- lassenianus* 192
- latus* 207, 215
- leiomerus* 208, 215
- sect. *Linearifolii* 123, 201, 219
- linearis* 186, 197, 199, 207, 215
- lobatus* 186, 188, 189, 206, 216
- macranthus* 217
- nauseosus* 204, 216
- multiceps* 189, 190
- neomexicanus* 359, 367
- nevadensis* 190, 191
 - var. *sonnei* 190
- nevadincola* 190, 191, 214
- ochroleucus* 197
 - var. *hirtellus* 197
- sect. *Oritrophium* 338
- ortegae* 187
- sect. *Osteocaulis* 199
- ovinus* 205, 216
- ozyphyllus* 187
- peregrinus* 186, 190, 204, 216, 217
 - var. *angustifolius* 190, 216
 - subsp. *callianthemus* 190

Erigeron peregrinus (cont.)var. *callianthemus* 190, 216var. *hirsutus* 190, 216, 217var. *scaposus* 190, 216*petiolaris* 187, 209*philadelphicus* 204, 217*pinkavii* 420*piperianus* 195-199*piscaticus* 188sect. *Polyactis* 200*pringlei* 201*pumilus* 186, 192, 193, 199, 200,
206, 212, 217subsp. *concinoides* 192, 212var. *gracilior* 217var. *intermedius* 192, 193, 217var. *pumilus* 192, 217*pygmaeus* 206, 217*radicatus* 199*scoparioides* 119, 122*scopulinus* 201*simplex* 187*sonnei* 190-192, 214*speciosus* 204, 217var. *macranthus* 217*tener* 208, 218sect. *Spathifolium* 199sect. *Trimorpha* 186, 203*uncialis* 207, 218subsp. *conjugans* 218var. *conjugans* 218var. *uncialis* 218*ursinus* 208, 218*utahensis* 209*vagus* 205, 218*velutipes* 188, 201*watsonii* 208, 219sect. *Wyomingia* 199

Eriocaulaceae 51

Eriocaulon 51*decangulare* 51*Eriochloa* 375*acuminata* 375*Eriochloa* (cont.)*gracilis* 375*lemmonii* 375

Eriogoneae 346

Eriogonoideae 345

Eriogonum 345, 346, 356, 372*abertianum* 372*deflexum* 372var. *turbinatum* 372*polycladon* 372*thurberi* 372*tomentosum* 345, 346*trichopes* 372*wrightii* 372*Erioneuron* 375*pulchellum* 375*Eriophyllum* 367*lanosum* 367*Erodium* 370*cicutarium* 370*texanum* 370*Eryngium* 51, 359, 366*heterophyllum* 359, 366*integrifolium* 51*Erysimum* 368*capitatum* 368*Erythrina* 359, 370*flabelliformis* 359, 370*Eschscholzia* 330, 331, 333, 336, 372*californica* 372ssp. *mezicana* 372*fumariifolia* 333*glyptosperma* 331*palmeri* 330*parishii* 331

Eschscholtzieae 330

Eschscholtzieae 330, 331

Eschscholtzioideae 331, 333

Eucalyptus 100*marginata* 100*Eucrypta* 371*micrantha* 371

- Eupatorieae 18, 23, 159, 304, 306,
 348
Eupatorium 51, 71, 359, 367
 fuertesii 71
 leucolepis 51
 perrinianum 71
 pycnocephalum 367
 rotundifolium 51
 solidaginifolium 359, 367
 tulanum 71
Euphorbia 370
 ezstipulata 370
 Euphorbiaceae 357, 366, 370
Eustachys 274-276
 caribaea 276
 distichophylla 274, 275
 floridana 274, 275
 glauca 274
 neglecta 274, 275
 petraea 274, 275
 retusa 274, 275
Euthamia 320
Evaz 367
 multicaulis 367
Evolvulus 369
 alsinoides 369
 arizonicus 369
 pilosus 369
 sericeus 369
Eysenhardtia 359, 370
 orthocarpa 370
 polystachya 359
 Fabaceae 124, 126, 281, 357, 370,
 425
 Fagaceae 370
Falcata 291
 ovalifolia 291
Ferocactus 369
 wislizenii 369
Filago 367
 californica 367
 depressa 367
Fimbristylis 125, 382, 383
 miliacea 382
Fleischmannia 21
 urenifolia 21
 var. *meziae* 21
Forestiera 371
 shrevei 371
Fouquieria 360, 370
 splendens 360, 370
 Fouquieriaceae 370
Frazinus 358, 371
 velutina 358, 371
 var. *toumeyii* 371
Froelichia 366
 arizonica 366
Fuirena 51
 squarrosa 51
 Fumariaceae 336

Galactia 285, 296, 359, 370
 hirta 296
 radicata 285
 wrightii 359, 370
Galinsoga 246, 255, 257
 sect. *Elata* 257
 filiformis 257
Galinsoginae 246, 255, 257, 258
Galium 373
 aparine 373
 microphyllum 373
 proliferum 373
Gaura 362, 372
 coccinea 372
 var. *arizonica* 372
 gracilis 372
 parviflora 372
Gayophytum 458
 strictum 458
Gelsemium 52
 sempervirens 52
 Gentianaceae 52
 Geraniaceae 370
 Gesneriaceae 220

- Gilia* 372
 mezicana 372
 ophthalmoides 372
 ssp. *australis* 372
Glandularia 374
 bipinnatifida 374
 var. *bipinnatifida* 374
Glinus 366
 radiatus 366
Glycine 289
 biloba 289
Gnaphalium 367
 chilense 367
 leucocephalum 367
 purpureum 367
 wrightii 367
Gomphrena 362, 366
 caespitosa 366
 sonorae 366
Gossypium 371
 thurberi 371
Gratiola 52
 neglecta 52
 pilosa 52
Grindelia 264-266, 326-329
 adenodonta 326, 327, 328
 arizonica 327
 grandiflora 327
 greenmanii 265
 inuloides 266
 lanceolata 327
 microcephala 326-328
 var. *adenodonta* 327
 var. *microcephala* 327, 328
 var. *montana* 326
 var. *pusilla* 327
 nuda 327
 oaxacana 326
 obovatifolia 265
 oxylepis 265, 327
 pusilla 326, 327, 328
 squarrosa 327
 tenella 327
Grindelia (cont.)
 turneri 264, 265
 villarrealii 264-266
Guilleminea 366
 densa 366
Guizotia 77, 80, 94
 bidentoides 94
Gutierrezia 363, 367
 microcephala 363, 367
 sarothrae 367
Gymnopogon 51
 brevifolius 51
Gymnospermae 366

Haloragaceae 370
Haplopappus 57
Haplophyton 359, 367
 crooksii 359, 367
Harpagonella 368
 palmeri 368
 var. *arizonica* 368
Hebecladus 274
 granulosus 273
Hedyotis 55
 uniflora 55
Helenieae 422
Helenium 358, 367
 thurberi 358, 367
Heliantheae 2, 13, 48, 96, 143, 148,
 149, 154, 242, 246, 255, 257,
 258, 302, 350, 422
Helianthus 51, 367
 angustifolius 51
 annuus 367
 petiolaris 367
Hemicarpha 374
 micrantha 374
Hemizonia 259, 260
 congesta 259, 260
 subsp. *congesta* 259
 leucocephala 259, 260
 var. *lutescens* 259
 corymbosa 260

Hemizonia (cont.)*leucocephala* 259, 260*lutescens* 259*multicaulis* 259 subsp. *vernalis* 259*Herissantia* 359, 371 *crispa* 359, 371*Heteranthera* 358, 376 *limosa* 358, 376*Heteropogon* 359, 360, 362, 375 *contortus* 360, 362, 375 *melanocarpus* 359, 375*Heterosperma* 154, 358, 367 *pinnatum* 358, 367*Heterotheca* 359, 365, 367, 416, 417,
449-455 *bolanderi* 450 *camphorata* 451 *echioides* 450 *fastigiata* 451 *horrida* 453 subsp. *cinerascens* 453 *rutteri* 359, 365, 367 *sessiliflora* 449-453 subsp. *bolanderi* 449, 451 subsp. *echioides* 449, 450, 451,
453 subsp. *fastigiata* 449, 451-453 var. *bolanderioides* 449, 450,
451 var. *camphorata* 449, 451,
453 var. *echioides* 450, 451 var. *fastigiata* 452, 453 var. *sanjacintensis* 449, 452 *subaxillaris* 367 *villosa* 449, 453, 454 var. *scabra* 449, 453, 454 var. *hispida* 453, 454 var. *shevockii* 449, 453, 454*Hibiscus* 371 *biseptus* 371 *coulteri* 371*Hilaria* 362, 375 *belangeri* 362, 375*Hofmeisteria* 18-23 *gayleana* 18-20, 22 *mexiae* 18, 21-23 *schaffneri* 20-22 *sinaloensis* 20-22 *standleyi* 21-23 *urenifolia* 18, 20, 21, 23 var. *mexiae* 18, 21*Hordeum* 359, 375 *murinum* 375 ssp. *glaucum* 375 *leporinum* 375 *pusillum* 359, 375*Hunnemannia* 330-336 *fumariaefolia* 336 *fumariifolia* 330-335 *hintoniorum* 330, 333-335*Hunnemannieae* 331*Hybanthus* 358, 374 *attenuatus* 358, 374 *verticillatus* 374*Hydastylus* 430*Hydrocotyle* 364, 366 *ranunculoides* 364, 366*Hydrophyllaceae* 61, 63, 358, 371*Hymenoclea* 367 *monogyra* 367*Hymenothrix* 367 *wislizenii* 367*Hypocoideae* 336*Hypericum* 52, 55 *brachyphyllum* 52, 55 *cruz-andreae* 52 *fasciculatum* 55 *hypericoides* 52*Hypocyrtia* 235*Hypoxis* 51 *rigida* 51*Ilex* 51, 178 *coriacea* 51

Ilex (cont.)*discolor* 178*opaca* 51*rubra* 178*vomitaria* 51*Ionactis* 416-418, 421-422*alpina* 417, 418, 421*caelestis* 418*elegans* 416, 420, 421*linariifolia* 417, 418*stenomeres* 418, 421*Ipomoea* 275, 356, 369*barbatisepala* 369*coccinea* 275, 369*costellata* 369*hederacea* 369*leptotoma* 369*purpurea* 369*quamoclit* 275*Ipomopsis* 372*longiflora* 372*Iresine* 359, 366*heterophylla* 359, 366

Iridaceae 374, 429, 430, 433, 434

Iridoideae 434

Isocoma 318, 360, 363, 367*coronopifolia* 318*tenuisecta* 360, 363, 367*Janusia* 371*gracilis* 371*Jatropha* 370*cardiophylla* 370*macrorrhiza* 370

Juglandaceae 371

Juglans 358, 371*major* 358, 371

Juncaceae 51, 374

Juncus 51, 363, 374*balticus* 363, 374*bufonius* 374*marginatus* 51*scirpoides* 51*Juncus* (cont.)*tenuis* 374var. *tenuis* 374*torreyi* 363, 374*trigonocarpus* 51*Justicia* 140-142sect. *Pentaloba* 140-142*hintoniorum* 140-142*pilosella* 140-142*turneri* 140-142*Kallstroemia* 374*californica* 374*grandiflora* 374*Karwinskia* 435-438*humboldtiana* 435, 438*tehuacana* 435, 436-438*Krameria* 360, 371*parvifolia* 360, 371var. *imparata* 371

Krameriaceae 371

Kyllinga 382*odorata* 382*pumila* 382*Lachnocaulon* 51*anceps* 51*Lactuca* 367*serriola* 367

Lemnaceae 374

Lactuceae 261

Laennecia 187, 201, 267-269*chihuahuana* 267, 268*confusa* 268*mapimiana* 269*pimana* 268*spellenbergii* 267, 268*Lagascea* 367*decipiens* 367

Lamiaceae 371

Lamium 371*amplexicaule* 371*Lappula* 368

- Lappula* (cont.)
 redowskii 368
Larrea 333, 354
 tridentata 354
Lasallea 422
Lasianthaea 24-26
 ritovegana 24-26
 rosei 24, 26
Lasthenia 367
 chrysostoma 367
Lauraceae 52
Leersia 364, 375
 oryzoides 364, 375
Leguminosae 155, 301, 428
Lemna 358, 364, 374
 gibba 364, 374
 minor 364, 374
 minima 358, 374
Lentibulariaceae 52
Lepidaploa 65, 67, 69, 75, 76
 buzifolia 69
Lepidium 368
 lasiocarpum 368
 var. *lasiocarpum* 368
 thurberi 368
 virginicum 368
 var. *medium* 368
Leptochloa 360, 375
 dubia 360, 375
 filiformis 375
 viscida 375
Leptoloma 375
 cognatum 375
Lesquerella 368
 gordoni 368
Lessingianthus 75
Leucelene 367, 417, 422
 ericoides 367, 417
Liatris 51
 pycnostachya 51
Liliaceae 51, 307, 374
Linaceae 52, 371
Linanthus 372
 Linanthus (cont.)
 aureus 372
Linaria 373
 tezana 373
Linum 52, 371
 lewisii 371
 medium 52
 puberulum 371
Loasaceae 371
Lobelia 52, 251, 253, 359, 369
 fenestralis 359, 369
 reverchonii 52
Loeflingia 369
 squarrosa 369
Loganiaceae 52
Lomatium 366
 nevadense 366
 var. *pseudorientale* 366
Lotus 357, 370
 corniculatus 370
 greenei 370
 humistratus 370
 oroboides 370
 salsuginosus 370
Ludwigia 372
 palustris 372
Lupinus 362, 370
 brevicaulis 362, 370
 concinus 370
 sparsiflorus 370
Lycium 373
 andersonii 373
 exsertum 373
Lycopodiaceae 51
Lycopodium 51
 alopecuroides 51
 appressum 51
 carolinianum 51
Lycurus 375
 setosus 375
Lythraceae 371
Lythrum 358, 371
 californicum 358, 371

- Machaeranthera* 318, 367, 368
 gracilis 367
 gypsophila 318
 pinnatifida 367
 ssp. *pinnatifida* 367
 restiformis 318
 tagetina 367
 tephrodes 368
Macroptilium 370
 heterophyllum 370
Macrosiphon 367
 brachysiphon 367
Madiinae 259, 260
Magnolia 52
 virginiana 52
Magnoliaceae 52
Malacothrix 368
 californica 368
 var. *glabrata* 368
 clevelandii 368
 fendleri 368
 sonchoides 368
Maloideae 135
Malpighiaceae 371
Malus 124, 130, 131
 pumila 124, 130
 sieboldii 124, 131
 sylvestris 124, 131
 toringo 131
Malva 371
 parviflora 371
Malvaceae 124, 126, 357, 371
Malvastrum 126, 127
 bicuspidatum 126, 127
 var. *glabrum* 127
 subsp. *tumidum* 126, 127
 var. *tumidum* 126, 127
Malvella 371
 leprosa 371
Mammillaria 359, 369
 gummifera 369
 var. *applanata* 369
 var. *macdougalii* 369
 Mammillaria (cont.)
 heyderi 359, 369
 var. *heyderi* 369
 macdougalii 369
 microcarpa 369
Manihot 370
 angustiloba 370
Marah 369
 gilensis 369
Marina 370
 calycosa 370
Marrubium 371
 vulgare 371
Marshallia 51
 graminifolia 51
 ssp. *tenuifolia* 51
Marsilea 358, 366
 vestita 358, 366
Marsileaceae 366
Martyniaceae 371
Matelea 277, 278, 280
 gonoloboides 280
 magallanesii 277, 278, 280
 pilosa 277, 280
 quirosii 280
Maurandya 373
 antirrhiniflora 373
Mecardonia 373
 vandellioides 373
Medicago 370
 polymorpha 370
Melampodium 368
 longicorne 368
Melastomataceae 52
Melilotus 370
 indicus 370
Menispermaceae 371
Mentzelia 371
 albicaulis 371
 asperula 371
 multiflora 371
 pumila 371
Mespilus 129, 130

- Mespilus* (cont.)
 arbutifolius 129, 130
Microlecanium 77, 80, 95, 96
 carinata 95
Microseris 368
 linearifolia 368
Microsteris 372
 gracilis 372
Mikania 159-163
 micrantha 164
 nana 159, 160
 obtusata 163
 obtusata 163
 pacei 159, 161, **163**
 reynoldsii 159, 162, **164**
 scandens 165
 viminea 159
Milleria 154
Mimosa 141, 360, 370
 biuncifera 360, 370
 dysocarpa 370
Mimulus 373
 floribundus 373
 guttatus 373
 nasutus 373
Minuria 340
Mirabilis 371
 longiflora 371
Mitracarpus 373
 breviflorus 373
Mollugo 366
 verticillata 366
Monactis 154
Monocotyledonae 374, 433
Monolepis 369
 nuttalliana 369
Monoptilon 416, 417
Montanoa 1
 speciosa 1
Moraceae 371
Morus 371
 microphylla 371
Muhlenbergia 51, 356, 363, 375
Muhlenbergia (cont.)
 arizonica 375
 asperifolia 363, 375
 emersleyi 375
 expansa 51
 fragilis 375
 microsperma 375
 porteri 375
 rigens 375
Myosurus 359, 373
 cupulatus 373
 minimus 359, 373
Myrica 52, 384
 cerifera 52
 heterophylla 52
Myricaceae 52
Myriophyllum 364, 370, 382
 aquaticum 382
 ezallescens 364, 370
Nama 61-63, 371
 cuatrocienegeense 61, 62
 hispidum 371
 var. *spathulatum* 371
 parvifolium 61, 62
 rzedowskii 61, 62
 serpylloides 61, 62
 var. *confertum* 62
 var. *serpylloides* 62
 var. *velutinum* 62
Nasturtium 368
 officinale 368
Neurocarpon 285
 mexicanum 285
Nicotiana 373
 trigonophylla 373
Nissolia 370
 schottii 370
Nolina 360, 374
 microcarpa 360, 374
Notholaena 359, 366
 grayi 366
 integerrima 366

Notholaena (cont.)*sinuata* 366*standleyi* 366*Nothoscordum* 363, 374*tezanum* 363, 374*Noticastrum* 339*Nyctaginaceae* 371*Nymphaea* 371*mezicana* 371*odorata* 371*Nymphaeaceae* 371*Nyssa* 52*sylvatica* 52*Nyssaceae* 52*Oberonia* 118*pusilla* 118*Octarrhena* 117, 118*pusilla* 117, 118*Odontotrichum* 40, 41, 46*eriocarpum* 46*pachyphyllum* 41*Oenothera* 357-359, 363, 372, 457-

459

sect. *Boisduvalia* 457*concinna* 457*densiflora* 457*flava* 359, 372*glabella* 459*primiveris* 372*pygmaea* 458*rosea* 358, 372*speciosa* 363, 372*subulata* 457*torreyi* 458*Oleaceae* 371*Olearia* 339, 340, 343*Onagraceae* 124, 127, 135, 357, 372,

456, 459, 460, 462

Oenothera 457, 458group *Boisduvalia* 457*cleistogama* 458*subulata* 457*Oenothera subulata* (cont.)race *concinna* 457*Opuntia* 356, 360, 369*arbuscula* 369*engelmannii* 369*fulgida* 369var. *mammillata* 369*leptocaulis* 369*macrorrhiza* 369var. *macrorrhiza* 369*phaeacantha* 369var. *discata* 369var. *major* 369*spinosior* 369*versicolor* 369*violacea* 369var. *santa-rita* 369*Orchidaceae* 3, 6, 51, 117, 124, 128,

443

Oritrophium 338-342*aciculifolium* 340*blepharophyllum* 338, 342*figueirasii* 338, 342*hirtopilosum* 340*nevadense* 338, 340, 342*orizabense* 338, 340-342*vahlhii* 338*Orobanchaceae* 372*Orobanche* 372*cooperi* 372*Orthocarpus* 373*purpurascens* 373*Osmunda* 51*cinnamomea* 51*regalis* 51*Osmundaceae* 51*Ostrya* 120*Oxalidaceae* 372*Oxalis* 372*albicans* 372ssp. *pilosa* 372*stricta* 372*Ozypolis* 51

- Ozypolis* (cont.)
rigidior 51
Pachystegia 339, 340
Panicum 51, 357, 375
antidotale 375
arizonicum 375
hirticaule 375
obtusum 375
pampinosum 375
stramineum 375
virgatum 51
Papaveraceae 330, 331, 336, 337, 372
Papavereae 331
Papaveroideae 336
Papilionateae 301
Paradrymonia 220, 231-233
binata 231
ciliosa 233
fuquaiana 232
hypocyrtia 232
prististoma 233
tylocalyca 232
Parietaria 373
hespera 373
Parkinsonia 358, 370
aculeata 358, 370
Parthenice 368
mollis 368
Paspalum 51, 55, 275, 364, 375
distichum 364, 375
floridanum 51, 55
laeve 55
notatum 275
Passiflora 359, 372
foetida 359, 372
mexicana 372
Passifloraceae 372
Pectis 368
longipes 368
Pectocarya 368
heterocarpa 368
platycarpa 368
Pectocarya (cont.)
recurvata 368
Pedicularaeae 414
Pellaea 359, 366
truncata 359, 366
Penstemon 373
parryi 373
Pentachaeta 418, 419, 423
bellidiflora 419
Pentadenia 220, 234-238
byrsina 235
ecuadorana 238
hypocyrtantha 234
isernii 236
katzensteinii 235
manabiana 235
rileyi 236
rubriacuta 237
tandapiana 238
trollia 234
zapotalana 236, 237
Perezia 73, 74
nervata 74
paniculata 73
vernonioides 73
Persea 52
borbonia 52
Perymenium 1, 2, 59, 60
hintonii 1, 2
paneroi 1, 2
Petromecon 330, 331
Petunia 373
parviflora 373
Phacelia 357, 358, 371
affinis 371
arizonica 371
coerulea 371
distans 371
ramosissima 358, 371
Phalaris 375
caroliniana 375
Phaseoleae 281
Phaseolus 359, 366

Phaseolus (cont.)*heterophyllus* 359*supinus* 366*Philodendron* 30, 32-38*alliodorum* 30, 32*ensifolium* 30, 32, 33*herbaceum* 30, 34, 35*hintoniorum* 59, 60*mendezii* 59, 60 *var. angustifolium* 59, 60*opacum* 30, 35-37 subg. *Pteromischum* 30*senatocarpium* 33*standleyi* 30, 37, 38*tamaulipense* 59, 60*yanezii* 59*Phoradendron* 374*californicum* 374*tomentosum* 374 ssp. *tomentosum* 374*Phreatia* 117, 118*baileyana* 117, 118*pusilla* 118*Phyla* 374*cuneifolia* 374*Physalis* 272, 373*hederaefolia* 373*longifolia* 373*orinocensis* 272*pubescens* 373*wrightii* 373

Phytolaccaceae 372

Picea 178*chiuahuana* 178*martinezii* 178

Pinaceae 51

Pinaropappus 261, 262*parvus* 261*pattersonii* 261, 262*pooleanus* 261, 262*Pinguicula* 52*pumila* 52*Pinus* 44, 51, 267, 427, 443*Pinus* (cont.)*cembroides* 427*palustris* 51, 443*taeda* 51*Piptocarpha* 68

Piptocarphinae 68

Plagiobothrys 368*arizonicus* 368*pringlei* 368

Plantaginaceae 372

Plantago 372*major* 372*patagonica* 372 *var. gnaphaloides* 372*virginica* 372*Platanthera* 443*integra* 443*Pleurophyllum* 339, 340, 343

Plumbaginaceae 372

Plumbago 359, 372*scandens* 359, 372*Poa* 363, 375*annua* 363, 375*bigelovii* 375*pratensis* 375

Poaceae 51, 274, 276, 356, 357, 375, 439

Podachaenium 143-148*andinum* 145*chiapanum* 143, 144-148*eminens* 143, 145*pachyphyllum* 143-146, 148*standleyi* 143-145, 146, 148

Podochilinae 117

Podochilus 117, 118*australiensis* 117, 118*Pogonia* 51*ophioglossoides* 51*Poinsettia* 370*heterophylla* 370 *var. graminifolia* 370 *var. heterophylla* 370*Polanisia* 369

Polanisia (cont.)*dodecandra* 369ssp. *trachysperma* 369*Polemoniaceae* 358, 372*Polygala* 52*incarnata* 52*Polygala* 372*barbeyana* 372*Polygalaceae* 52, 372*Polygonaceae* 345, 346, 357, 372*Polygonum* 357, 358, 364, 372*aviculare* 372*lapathifolium* 372*pensylvanicum* 358, 372*persicaria* 372*punctatum* 364, 372*Polypogon* 363, 364, 375*monspeliensis* 363, 364, 375*Pontederiaceae* 376*Populus* 358, 373*fremontii* 358, 373*Porophyllum* 368*gracile* 368*ruderales* 368ssp. *macrocephalum* 368*Portulaca* 362, 372*retusa* 372*suffrutescens* 372*umbraticola* 372*Portulacaceae* 372*Potamogeton* 358, 376*foliosus* 358, 376*pusillus* 358, 376*Potamogetonaceae* 376*Potentilla* 359, 373*rivalis* 359, 373*Primulaceae* 372*Proboscidea* 371*altheaefolia* 371*parviflora* 371*Prosopis* 333, 358, 360, 370*velutina* 358, 360, 370*Psacalium* 40-48*Psacalium* (cont.)*amplum* 41, 42, 44*cronquistiorum* 40, 42-45*eriocarpum* 40, 45-47*pachyphyllum* 40-45*pentaflorum* 40, 45-47*sinuatum* 44*Pteridium* 51*aquilinum* 51*Pteridophyta* 366*Pteromischum* 33*Ptilimnium* 51*capillaceum* 51*Pyrrhopappus* 363, 368*multicaulis* 368*rothrockii* 363, 368*Pyrrocoma* 57, 58*carthamoides* 57var. *subsquarrosa* 57*clementis* 57var. *villosa* 57*congesta* 58*crocea* 58var. *genuflexa* 58*genuflexa* 58*hirta* 58var. *lanulosa* 58*lanceolata* 58var. *subviscosa* 58*lanulosa* 58*racemosa* 58var. *congesta* 58var. *sessiliflora* 58*sessiliflora* 58*subsquarrosa* 57*subviscosa* 58*villosa* 57*Pyrus* 130, 131*japonica* 130*malus* 130, 131var. *paradisiaca* 130var. *sylvestris* 131*sieboldii* 131

Quercus 120, 137, 178, 267, 268, 358-360, 370

coccolobifolia 268

eduardii 137

emoryi 358, 360, 370

cf. *hartwegii* 267

macvaughii 268

oblongifolia 358, 360, 370

scytophylla 137

turbinella 359, 360, 370

viminea 137

Rafinesquia 368

californica 368

neomezicana 368

Ranunculus 363, 364, 373

hydrocharoides 364, 373

macranthus 363, 373

Remya 340, 344

Rhamnaceae 373, 435, 438

Rhezia 52

mariana 52

petiolata 52

Rhododendron 52

canescens 52

Rhoogeton 220, 239

panamensis 239

viviparum 239

Rhynchosia 370

tezana 370

Rhynchosida 371

physocalyz 371

Rhynchospora 51

chalarocephala 51

globularis 51

var. *globularis* 51

gracilentia 51

macra 51

oligantha 51

plumosa 51

pusilla 51

Richardia 275

brasiliensis 275

Rigiopappus 418

Rivina 372

humilis 372

Rosaceae 124, 129, 135, 373

Rubiaceae 373

Rubus 275

Rumex 372

crispus 372

hymenosepalus 372

Sabal 16

Sabatia 52

gentianoides 52

Sabazia 246, 257

Sacciolepis 382

indica 382

Sageretia 373

wrightii 373

Salicaceae 373

Salix 358, 363, 373, 460

cf. *babylonica* 373

gooddingii 358, 363, 373

taxifolia 358, 373

Salsola 369

iberica 369

Sambucus 369

mxicana 369

Sapindaceae 373

Sapindus 358, 373

saponaria 358, 373

var. *drummondii* 358, 373

Sapotaceae 373

Saracha 378, 380

biflora 378, 380

Sarcostemma 367

crispum 367

cynanchoides 367

ssp. *cynanchoides* 367

ssp. *hartwegii* 367

Sarracenia 52

alata 52

Sarraceniaceae 52

Saururaceae 373

- Schismus* 375
 barbatus 375
Schizachyrium 51, 55, 359, 375
 cirratum 359, 375
 scoparium 51, 55
 tenerum 51, 55
Scirpus 125, 358, 364, 374
 californicus 358, 374
 confervoides 125
 olneyi 364, 374
Scleria 51, 267
 ciliata 51
 georgiana 51
 reticularis 51
Scrophulariaceae 52, 137, 139, 247,
 250, 251, 357, 373, 384, 388,
 389, 414, 415
Senecio 48, 51, 359, 368
 douglasii 368
 var. *douglasii* 368
 var. *longilobus* 368
 lemmoni 359, 368
 tomentosus 51
Senecioneae 40, 48
Senna 124, 126, 275, 370
 artemisioides 124, 126
 bauhinioides 370
 hirsuta 370
Setaria 363, 375
 arizonica 375
 geniculata 363, 375
 grisebachii 375
 leucopila 375
Sicyosperma 358, 369
 gracile 358, 369
Sida 371
 procumbens 371
 spinosa 371
 var. *angustifolia* 371
Sidalcea 124, 126, 127, 134, 363, 371
 malviflora 124, 126, 127
 subsp. *laciniata* 124, 126, 127
 var. *laciniata* 124, 126, 127
Sidalcea malviflora (cont.)
 var. *sancta* 126, 127
 neomexicana 363, 371
Sideroxylon 358, 373
 lanuginosa 358, 373
Silene 369
 antirrhina 369
Simmondsia 358, 373
 chinensis 358, 373
Simmondsiaceae 373
Siphonoglossa 366
 longiflora 366
Sisymbrium 368
 irio 368
Sisyrrinchieae 434
Sisyrrinchium 363, 374, 429-434
 sect. *Bermudiana* 433
 biforme 429, 432
 conzattii 433
 demissum 363, 374, 429, 432, 433
 dimorphum 429, 432-434
 subg. *Echthronema* 430-432, 434
 ensigerum 432
 micranthum 429, 432, 433
 novoleonense 429, 430-432
 quadrangulatum 431
 scabrum 429, 430, 432, 433
 sect. *Sisyrrinchium* 430
 subgen. *Sisyrrinchium* 429-432
Siphonoglossa 140, 142
Sitanion 375
 hystrix 375
Smilax 51
 laurifolia 51
Smithiantha 220, 239, 240
 canarina 239
 loui 240
 multiflora 240
Solanaceae 166, 180, 183, 270, 357,
 373, 378
Solanum 166-168, 180-185, 373, 378-
 380
 acroglossum 180

Solanum (cont.)

- albornozii* 180
- amabile* 166
- amayanum* 180
- anamatophilum* 180
- antacochense* 180
- ayacuchense* 180
- blanco-galdosii* 180
- brevicaule* 185
- buesii* 180
- bukasovii* 185, 166, 167, 380
 - var. *multidissectum* 167
- cacetanum* 167
- canasense* 166, 167
 - var. *alba* 166
 - var. *calcense* 166
 - var. *intihuatanense* 166
 - var. *latifolia* 166
 - var. *lechnoviczii* 166
 - var. *xerophilum* 167
 - var. *xerophylla* 167
- candolleianum* 181
- cantense* 181
- colombianum* 167
- series *Conicibaccata* 185
- contumazaense* 181
- chilliasense* 181
- chiquidenum* 181
- deflexum* 373
- dolichocremastrum* 181
- douglasii* 373
- elaeagnifolium* 373
- espinarense* 167
- guzmanguense* 181
- hapalosum* 167
- hawkesii* 168
- humectophilum* 181
- hypacrarthrum* 181
- immitte* 181
- ingaefolium* 181
- irosinum* 181
- jalcae* 181
- lechnoviczii* 167

Solanum lechnoviczii (cont.)

- var. *latifolia* 167
 - var. *xerophylla* 167
 - litusinum* 181
 - lobbianum* 181, 183-185
 - f. *multidissectum* 185
 - lumholtzianum* 373
 - marinasense* 167
 - minutifolium* 181
 - multidissectum* 167
 - nemorosum* 181
 - neovavilovii* 181
 - nubicola* 182
 - ochoae* 167
 - olmosense* 182
 - orophilum* 182
 - pampasense* 167, 182
 - paramoense* 168, 182
 - peloquinianum* 182
 - sect. *Petota* 166, 378
 - pillahuatense* 182
 - piurae* 182
 - pumilum* 167
 - punoense* 167
 - quillonanum* 182
 - raphanifolium* 168
 - raquialatum* 182
 - saxatilis* 378-380
 - sicuanum* 167
 - soukupii* 167
 - var. *espinarense* 167
 - suffrutescens* 182
 - tacnaense* 182
 - taulisense* 182
 - trinitense* 182
 - series *Tuberosa* 380
 - tuberosum* 168
 - subsp. *andigena* 168
 - urubambae* 182
 - villuspetalum* 182
 - wittmackii* 182
- Solidago* 368, 422
- sparsiflora* 368

- Sonchus* 368
 asper 368
Sorghum 360, 375
 halepense 360, 375
Spermolepis 366
 echinata 366
Sphaeralcea 358, 371
 angustifolia 371
 emoryi 371
 fendleri 358, 371
 var. *venusta* 371
Sphagnum 50, 509
Spiranthes 124, 128
 confusa 124, 128
 durangensis 128
Spiranthinae 134
Sporobolus 363, 375, 376
 contractus 375
 cryptandrus 375
 wrightii 363, 375, 376
Squamopappus 143, 148
Stachys 371
 coccinea 371
Stemodia 251-254
 costaricensis 251-253
 reliquiarum 251-253
Stenocarpa 255-258
 filiformis 255
 ritovegana 255, 256
Stenocephalum 75
Stephanomeria 368
 ezigua 368
 pauciflora 368
Sterculiaceae 373
Stipa 378, 380
 ichu 378, 380

Talinum 372
 aurantiacum 372
 paniculatum 372
Tamaricaceae 373
Tamarix 373
 ramosissima 373

Taxus 178
 globosa 178
Tecoma 359, 368
 stans 359, 368
Tephrosia 370
 tenella 370
Tetraclea 374
 coulteri 374
Tetramalopium 340
Tetramerium 366
 nervosum 366
 hispidum 366
Thelasiinae 117
Thelypodopsis 358, 368
 linearifolia 358, 368
Thysanocarpus 368
 curvipes 368
 var. *elegans* 368
Tidestromia 366
 lanuginosa 366
Tilia 120
Tillaea 369
 erecta 369
Tithonia 358, 368
 thurberi 358, 368
Toxicodendron 51
 radicans 51
 vernix 51
Trachypogon 359, 376
 secundus 359, 376
Tracyina 418, 421
Tragia 370
 nepetaefolia 370
Trianthema 366
 portulacastrum 366
Tribulus 374
 terrestris 374
Trichachne 375
 californica 375
insularis 375
Tridax 243, 246, 350-352
 candidissima 350, 352
 hintoniorum 350-352

Tridens 51*ambiguus* 51*Trifolium* 363, 370*wormskioldii* 363, 370var. *wormskioldii* 370*lacerum* 370*Trimorpha* 186, 201, 203, 204, 208*lonchophyllus* 203, 204, 208*Triodanis* 369*biflora* 369*holzingeri* 369*Trisetum* 376*interruptum* 376*Triticum* 376*aestivum* 376*Trizix* 368*californica* 368*Typha* 364, 376*domingensis* 376*latifolia* 364, 376

Typhaceae 376

Ulmaceae 373

Urticaceae 373

Utricularia 52, 441*cornuta* 52*junceae* 52*subulata* 52*Vaccinium* 52*corymbosum* 52*Verbena* 321-325, 359, 374*bracteata* 359, 374*canescens* 321, 323*gracilis* 321, 374*halei* 321*johnstonii* 321, 323, 324*menthaefolia* 321*neomexicana* 321, 323, 374var. *hirtella* 321, 323var. *neomexicana* 323*perennis* 321-324forma *johnstonii* 322*Verbena perennis* (cont.)var. *johnstonii* 321*plicata* 321*shrevei* 322*stricta* 325

Verbenaceae 321, 325, 374

Verbesina 14, 15, 145, 154, 302, 303, 368*encelioides* 368*hintoniorum* 302, 303*montanoifolia* 14, 15*potosina* 303sect. *Pseudomontanoa* 14, 15*standleyi* 145*tecolotlana* 14*zaragosana* 302, 303*Vernonanthura* 65, 66-74*almedae* 68*amplexicaulis* 66, 68*angulata* 66, 68*auriculata* 69*beyrichii* 69*brasiliiana* 69*buxifolia* 69*canaminina* 69*catharinensis* 69*chamaedrys* 69*cichoriifolia* 69*chaquensis* 69*cocleana* 66, 69*condensata* 69*cordata* 70*crassa* 66, 70*cronquistii* 70*cuneifolia* 70*cupularis* 70*cymosa* 68, 70*deppeana* 70*diffusa* 70*discolor* 70*fagifolia* 70*ferruginea* 70*fuertesii* 71

Vernonanthura (cont.)

- havanensis* 71
- hieracioides* 71
- ignobilis* 71
- laxa* 71
- liatroides* 71
- loretensis* 71
- lucida* 71
- mariana* 71
- membranacea* 68, 71
- menthaefolia* 71
- montevidensis* 68, 72
- mucronulata* 72
- nudiflora* 72
- oaxacana* 72
- oligactoides* 72
- oligolepis* 72
- paludosa* 72
- patens* 72
- petiolaris* 73
- phaeoneura* 73
- phosphorica* 68, 73
- pinguis* 66, 73
- piresii* 73
- prenanthoides* 73
- puberula* 73
- rigiophylla* 73
- sambrayana* 73
- serratuloides* 73
- sinclairii* 74
- squamulosa* 74
- stellata* 74
- subverticillata* 74
- tuerckheimii* 74
- tweedieana* 66, 74
- viscidula* 74
- westiana* 74
- yurimaguasensis* 74

Vernonia 16, 17, 65, 67-76

- albifila* 74
- almedae* 68
- amplexicaulis* 68
- angulata* 69

Vernonia (cont.)

- arechavaletae* 72
- aschenborniana* 72
- assana* 69
- auriculata* 69
- baccharoides* 72
- bahiensis* 70
- bangii* 72
- barkeri* 71
- beyrichii* 69
- blodgettii* 67
- brasiliansa* 68, 69
- ser. *Brasilianae* 67
- buzifolia* 69
- subsect. *Buzifoliae* 67
- camporum* 74
- canaminina* 69
- capreaefolia* 71
- catharinensis* 69
- chamaedrys* 69
- subsect. *Chamaedrys* 67, 76
- subser. *Chamaedrys* 67
- subser. *Chaquensis* 67
- cichoriifolia* 69
- chaquensis* 69
- cocleana* 69
- collina* 72
- condensata* 69
- conyzoides* 72
- corcovadensis* 73
- cordata* 70
- corymbulosa* 74
- crassa* 70
- crenata* 70
- cronquistii* 70
- cubensis* 71
- cuneifolia* 70
- cupularis* 70
- cymosa* 70
- denticulata* 69
- deppeana* 70
- diffusa* 70
- discolor* 70

Vernonia (cont.)

- domingensis* 69
- ehrenbergiana* 71
- elsiae* 73
- ervendbergii* 67
- fagifolia* 70
- faustiana* 16, 67
- ferruginea* 70
- florida* 72
- fuertesii* 71
- gochnatioides* 72
- greggii* 67
- grisebachii* 71
- haenkeana* 72
- havanensis* 71
- hebeclada* 74
- hieracioides* 71
- hilariana* 73
- ignobilis* 71
- ilex* 69
- insularis* 67
- itapensis* 70
- jaliscana* 73
- joyaliae* 16
- karvinskiana* 74
- lanceolaris* 72
- laxa* 71
- subser. *Laxae* 67
- sect. *Lepidaploa* 16, 17
- subsect. *Lepidaploae Paniculatae* 67
- subsect. *Lepidaploae Scorpioideae* 67
- liatroides* 71
- loretensis* 71
- lucida* 71
- mariana* 71
- membranacea* 71
- menthaefolia* 71
- micradenia* 72
- microphylla* 74
- missionis* 68, 70
- monsonensis* 72

Vernonia (cont.)

- mucronulata* 72
- montana* 69
- montevidensis* 72
- morelana* 70
- nitidula* 68, 72
- ser. *Nitidulae* 67
- subsect. *Nitidulae* 67
- nudiflora* 72
- subsect. *Nudiflorae* 67
- subser. *Nudiflorae* 67
- oazacana* 72
- odoratissima* 69
- oligactoides* 72
- oligolepis* 72
- orientis* 71
- ottonis* 71
- pacchensis* 72
- paludosa* 72
- subsect. *Paniculatae* 67, 75
- patens* 72
- petiolaris* 73
- phaeoneura* 73
- pinguis* 73
- piresii* 73
- polyanthes* 68, 73
- subsect. *Polyanthes* 67
- polycephala* 70
- praecox* 74
- prenanthoides* 73
- psittacorum* 73
- puberula* 73
- ser. *Puberulae* 67
- pusilliflora* 74
- rigiophylla* 73
- ruficoma* 68, 71
- salamana* 72
- sambrayana* 73
- scabra* 69
- schaffneri* 67
- schulziana* 74
- subsect. *Scorpioidea Verae* 67
- serratuloides* 73

Vernonia (cont.)

- sinclairii* 74
- sorocabae* 72
- squamulosa* 74
- stictophylla* 71
- stuebelii* 72
- subverticillata* 74
- sylvestris* 70
- tuerckheimii* 74
- tweddieana* 74
- umbellifera* 73
- ser. *Umbelliformes* 67, 74, 75
- vargasii* 72
- vernonioides* 74
- viscidula* 74
- weberbaueri* 72
- westiana* 74
- yurimaguasensis* 74

Vernoniaceae 75*Vernonieae* 17, 65, 66, 75, 76*Vernoniinae* 65, 66*Veronica* 359, 373

- anagallis-aquatica* 373
- peregrina* 359, 373
- ssp. *zalapensis* 359, 373

Viburnum 52

- nudum* 52

Vicia 370

- ludoviciana* 370

Viguiera 1, 368

- dentata* 368
- var. *lancifolia* 368
- multiflora* 368
- puruana* 1

Viola 52, 509

- primulifolia* 52

Violaceae 52, 374, 509*Viscaceae* 374*Vitaceae* 124, 132, 374*Vitex* 16

- mollis* 16

Vitis 132, 133, 358, 374

- aestivalis* 133

Vitis aestivalis (cont.)

- var. *bourquiniana* 133
- var. *glauca* 133
- arizonica* 358, 374
- var. *arizonica* 374
- candicans* 132, 133
- var. *coriacea* 132, 133
- cinerea* 132, 133
- var. *canescens* 132, 133
- cordifolia* 132, 133
- var. *helleri* 132, 133
- coriacea* 132
- longii* 132, 133
- var. *microsperma* 132, 133
- rupestris* 132, 133
- var. *dissecta* 132, 133
- vulpina* 133
- var. *praecox* 133

Vulpia 376

- octoflora* 376

Vittadinia 340*Waltheria* 359, 373

- americana* 359, 373

Websteria 124, 125

- confervoides* 124, 125

Wedelia 48*Xanthium* 363, 368

- strumarium* 363, 368

Xylothamia 318-320

- pseudobaccharis* 319
- purpusii* 319
- triantha* 318, 319
- truncata* 318, 319

Xyridaceae 51*Xyris* 51, 55, 441

- ambigua* 51
- baldwiniana* 51
- difformis* 51
- var. *curtissii* 51
- drummondii* 51, 55
- jupicai* 55

Xyris (cont.)

scabrifolia 51

torta 51

Yabea 366

microcarpa 366

Yucca 120, 333, 360, 374

arizonica 374

elata 360, 374

Zamia 141

Zannichellia 364, 376

palustris 364, 376

Zannichelliaceae 376

Zephyranthes 374

longifolia 374

Zigadenus 307, 308, 311

fontanus 308

micranthus 307, 308

var. *fontanus* 307, 308

var. *micranthus* 308

Zinnia 360, 368

acerosa 360, 368

Zizyphus 363, 373

obtusifolia 363, 373

var. *canescens* 373

Zygophyllaceae 374

CORRECTIONS AND ADDITIONS

Volume 71, number 6, inside front cover, add to Contents:

"Index to authors, volume 71.	486
Index to taxa, volume 71.	487
Index to reviewers, volume 71. dotfill	510
Disposition of manuscripts received for volumes 70 and 71.	512"

Volume 72, number 1, page 37, line 5 of the Introduction, replace 'species.' This' with 'species'. This'.

Volume 72, number 1, page 38, line 1, replace "Replaced synonym" with "REPLACED SYNONYM".

Volume 72, number 1, page 38, line 28, replace "LECTOTYPE [here" with "LECTOTYPE NOV. [here".

Volume 72, number 2, cover, L.E. MCKINNEY is the second author for the paper entitled "A nomenclatural change in *Viola* (Violaceae)

Volume 72, number 2, page 99, line 3 of abstract, replace "(= *Coreopsis scopulorum*)" with "(= *Coreopsis scopulorum* Sherff)".

Volume 72, number 2, page 99, line 5 of abstract, replace "(= *Coreopsis morotonensis*)" with "(= *Coreopsis morotonensis* Sherff)".

Volume 72, number 2, page 99, line 6 of abstract, replace "(= *Coreopsis odora*)" with "(= *Coreopsis odora* Sherff)".

Volume 72, number 2, page 99, line 7 of abstract, replace "(= *Bidens paupercula* var. *filirostris*)" with "(= *Bidens paupercula* Sherff var. *filirostris* P. Taylor)".

Volume 73, number 1, page 50, replace "*Sphagnum*" with "*Sphagnum*."

Volume 73, number 3, page 198, see detailed statement on page 463 of volume 73, number 6.

INDEX TO REVIEWERS, VOLUME 73

The editor express his most sincere appreciation to the following individuals. These are persons who have reviewed papers that were submitted for publication in volume 73 of *Phytologia*. Without the willingness and diligence of these reviewers, the task of the editor would be much more difficult, and the quality of the papers published would be lessened. To each of you, I offer my most sincere thanks.

Michael J. Warnock, Editor.

Baldwin, B.G.
Barneby, R.
Barrie, F.R..
Berry, P.
Boufford, D.
Boyd, S.
Breedlove, D.
Brothers, L.
Brown, L.E.
Brummitt, R.K.
Canne-Hilliker, J.
Capels, K.
Carter, R.
Clark, W.D.
Clements, M.A.
Cronquist, A.
Crosby, M.
Crawford, D.
Dodd III, T.
Dorn, B.
Dressler, R.L.
Dutton, B.
Ertter, B.
Escobar, L.A.

Felger, R.
Funk, V.A.
Grant, J.
Gutknecht, K.
Hartman, R.
Holmgren, N.
Jacobs, S.
Jansen, R.
Jarvis, C.E.
Jeffrey, C.
Jones, G.D.
Jury, S.L.
Kartesz, J.
Kral, R.
Lellinger, D.B.
Lievens, A.
Lonard, R.I.
de la Luz Arreguín, M.
MacRoberts, D.T.
Martínez, M.
Mason, C.
Mayfield, M.
Maze, J.
McDaniel, S.

McVaugh, R.	Solomon, J.
Moore, D.M.	Soule, J.
Nesom, G.L.	Spellenberg, R.
Nicolson, D.H.	Strong, M.
Panero, J.	Sundell, E.
Peterson, P.	Terrell, E.E.
Pinkava, D.J.	Thorne, R.F.
Powell, A.M.	Todzia, C.A.
Pruski, J.	Townsend, A.M.
Raj, N.	Turner, B.L.
Ramamoorthy, T.P.	Wallace, B.J.
Reveal, J.L.	van der Werff, H.
Riggins, R.L.	Wiersma, J.
Robinson, H.	Wilken, D.
Rodríguez, C.	Williams, N.A.
Sanders, A.C.	Wipff, J.K.
Schatz, G.E.	Wurdack, J.
Smith, S.	

DISPOSITION OF MANUSCRIPTS RECEIVED FOR VOLUMES 72 AND 73

Manuscripts listed below include those received after examination by two or more reviewers. A note of appreciation to reviewers is found on page 510 with the list of reviewers. Manuscripts received without review are not considered for publication until review has been completed.

Manuscripts received: 135

Manuscripts accepted without revision: 8

Manuscripts accepted after revision: 113

Manuscripts returned to authors without publication: 7

Manuscripts currently under revision: 7


Manuscripts currently accepted but not yet published: 5

Papers published: 116

Days elapsed from receipt of manuscript to publication (includes only those manuscripts selected for publication): mean = 45; range = 22-141

Days elapsed from acceptance for publication to publication of manuscript:
mean = 28; range = 15-51

New York Botanical Garden Library



3 5185 00288 4664

Information for Authors

Articles from botanical systematics and ecology, including biographical sketches, critical reviews, and summaries of literature will be considered for publication in PHYTOLOGIA. Manuscripts may be submitted either on computer diskette, or as typescript. Diskettes will be returned to authors after action has been taken on the manuscript. Diskettes may be 5.25 inches or 3.5 inches and may be written in any IBM or MacIntosh compatible format. Typescript manuscripts should be single spaced and will be read into the computer using a page scanner. The scanner will read standard typewriter fonts but will not read dot matrix print. Manuscripts submitted in dot matrix print cannot be accepted. Use underscore (not italics) for scientific names. Corrections made on typescript manuscripts must be complete and neat as the scanner will not read them otherwise. Language of manuscripts may be either English or Spanish. Figures will be reduced to fit within limits of text pages and therefore, should be submitted with an internal scale and have dimensions proportional to those for text pages. Legends for figures should be included in figures whenever possible. Each manuscript should have an abstract and key word list. Specimen citations should be consistent throughout the manuscript. Serial titles should be cited with abbreviations used in *Botanico Periodicum Huntianum*. References cited only as part of nomenclatural summaries should not appear in Literature Cited. Nomenclatural work should include one paragraph per basionym and must provide proper (as defined by the current *International Code of Botanical Nomenclature*) citation of sources of epithets and combinations.

Authors should arrange for two workers in the appropriate field to review the manuscript before submission. Copies of reviews should be forwarded to the editor with the manuscript. Manuscripts will not be published without review.

Cost of publication is currently \$13.00 US per page for publication without reprints. Publication with 100 reprints is provided for \$18.00 US per page, 200 reprints for \$21.50 US per page. Page charges are due with manuscript and no paper will be published before payment is received in full. Reprints must be ordered and paid for in advance. Page charges will be determined on the basis of a typescript page (single spaced, 10 points, blank line between paragraphs) with all type inside a rectangle 143 mm (horizontal) by 219 mm (vertical), not including running head and page number. Title page should include title, author(s) name(s), and address(es). Two blank lines should appear above and below section headings (Abstract, Discussion, Literature Cited, etc.) in the manuscript. No extra charge is made for line drawings provided they conform to limitations of size and proportion for normal text. Halftones require an extra charge of \$10.00 US per page.